



James Cooper Lane

Philadel

MS 2/0059-01

Acc. 1987-013 (1)

Handwritten notes in cursive script, possibly mentioning "Society" and "Providence".

cta. D. D.

John Wiley Book

October 15th 1835

CARD 500
20-0410
LOCATION

Handwritten notes and a circular stamp at the bottom left, possibly containing the word "Church".

Wiley *John Wiley*
33 Walnut St
1867

MEMORIA MEDICA.

A

MEDICAL COMMON PLACE BOOK,

WITH AN

ALPHABETICAL INDEX

OF THE

MOST COMMON TERMS OCCURRING IN PRACTICE.

John Wiley

CAREFULLY SELECTED AND ARRANGED BY
A FELLOW OF THE MASSACHUSETTS MEDICAL SOCIETY.

BOSTON:
PUBLISHED BY CARTER & HENDEE.

MDCCCXXIX.

DISTRICT OF MASSACHUSETTS.....TO WIT:

District Clerk's Office.

Be it remembered, that on the fourteenth day of September, A. D. 1829, in the fifty fourth year of the Independence of the United States of America, CARTER & HENDEE, of the said District, have deposited in this Office the title of a Book, the right whereof they claim as Proprietors, in the words following, *to wit* :

"Memoria Medica. A Medical Common Place Book, with an Alphabetical Index of the most common terms occurring in practice. Carefully selected and arranged by a Fellow of the Massachusetts Medical Society."

In conformity to the Act of the Congress of the United States, entitled "An Act for the encouragement of learning, by securing the copies of maps, charts, and books, to the authors and proprietors of such copies, during the times therein mentioned:" and also to an Act entitled "An Act supplementary to an Act entitled An Act for the encouragement of learning, by securing the copies of maps, charts, and books, to the authors and proprietors of such copies, during the times therein mentioned; and extending the benefits thereof to the arts of designing, engraving and etching historical and other prints."

JNO. W. DAVIS, { *Clerk of the District
of Massachusetts.*

PREFACE.

John Wiley

MEDICAL knowledge is of all kinds the most fugacious and difficult to be fixed. The immense multiplicity of facts which crowd the memory of the experienced practitioner, defies the most retentive powers. It has been correctly observed, that a Physician never makes use of more than ten years' experience, and beyond that, *forgets* as much as he *learns*. A common place book supplies in a considerable degree this defect so universally felt and complained of. Every medical man, whether pupil or practitioner, is constantly meeting with some fact or observation, which it would be of use to him hereafter to recur to, and yet it would be in vain to trust the recollection of it to his memory alone. This is peculiarly true of successful extemporaneous prescriptions, and of curious cases and unexpected recoveries, and facts disclosed by the examination of morbid parts after death.

The following work is intended to furnish a convenient depository of this fleeting and ephemeral knowledge; and it is hoped that it will be made useful, not only to practitioners themselves, but likewise to the public, by enabling any one to communicate the results of his experience with comparatively little trouble. A large list of the most important terms occurring in general practice, is alphabetically arranged, and a convenient space is left for the addition of new terms. In order not to swell this list, the terms used in Chemistry have been left out, as their number would require a common place book by itself.

The mode of using a common place book upon this plan, is commonly known. Suppose a case of cutaneous disease, which has long baffled the attempts to cure it, suddenly yields to a new and fortunate prescription. These facts are recorded, and the number of the page in which the record is placed, is put against the words "cutaneous diseases." The same number may likewise, without the least confusion, be placed against several other words, as "skin," "prescription," "cure." In this way every thing which is recorded may be found without trouble.

OCTOBER 1, 1829.

PREFACE

Medical knowledge is of all kinds the most fugitive and difficult to be fixed. The immense multiplicity of facts which crowd the memory of the experienced practitioner, defies the most retentive power. It has been correctly observed, that a Physician never takes use of more than two or three experiments, and beyond that, as much as he knows. A common-place book supplies a considerable defect in this respect so universally felt, and recommended. Every practical man, who would be at all proficient, is constantly receiving with some fact or observation, which it would be useless to him to retain to recur to, and yet it would be in vain to trust the recollection of it to his memory alone. This is peculiarly true of surgery, and of the numerous principles, and of cases and unexpected reactions, and their effects by the examination of wounds, parts after death.

The following work is intended to furnish a convenient repository of this fugitive and ephemeral knowledge; and it is hoped that it will be made useful, not only to practitioners themselves, but likewise to the public, by enabling them to be conversant with the results of the experience with comparatively little trouble. A large part of the most important terms occurring in general practice, is alphabetically arranged, and a convenient space is left for the addition of new terms. In order not to swell this list, the terms used in Chemistry have been left out, as their number would require a common-place book by itself.

The mode of using a common-place book upon this plan is commonly known. Suppose a case of catarrhus vesicæ, which has long baffled the attempts to cure it, suddenly yields to a new and fortunate prescription. These facts are recorded, and the number of the page in which the record is placed is put against the words "catarrhus vesicæ." The same number may likewise be put against the words "cure." In this way, every thing which is recorded may be found without trouble.

October 31, 1811.

Dr. Colhoun's Lectures

S. M. Clellan's

Rever's

Pattison's

Green's

S. M. Clellan's

Dunglison's

57
75
154

143
169

67

Notes taken from

INDEX.

	Page.		Page.
ABDOMEN,		ANTACIDS,	
ABORTION,		ANTHELMINTICS,	
ABSCCESS,		ANTHRAX,	
ABSORBENTS,		ANTIDOTES,	
ABSORPTION,		ANTIMONY,	
ADHESION,		ANTISEPTICS,	
ACUPUNCTURATION,		ANTISPASMODICS,	
AFTER PAIN,		ANUS,	
AGUE,		ANXIETY,	
AIR,		AORTA,	
ALCALIES,		APERIENTS,	
ALCOHOL,		APHONIA,	
ALIMENTARY CANAL,		APOPLEXY,	
ALOES,		APPETITE,	
ALTERATIVES,		APTHÆ,	
AMAUROSIS,		AROMATICS,	
AMENORRHÆA,		ARSENIC,	
AMMONIACUM,		ARTERY,	
AMMONIA,		ARTHRITIS,	
AMPUTATION,		ARTICULATION,	
ANASARCA,		ASCARIDES,	
ANASTOMOSIS,		ASCITES,	
ANATOMY,		ASPHYXIA,	
ANCHYLOSIS,		ASSAFÆTIDA,	
ANEURISM,		ASTHMA,	
ANGINA PECTORIS,		ASTRINGENTS,	
ANODYNES,		ATROPHIA,	
ANOREXIA,		AUSCULTATION,	

Page.

BALSAM,
 BANDAGES,
 BARK,
 BARRENNESS,
 BARYTES,
 BATH,
 BATTERY, GALVANIC,
 BED,
 BEET,
 BELLADONNA,
 BILE,
 BISMUTH,
 BLADDER,
 BLINDNESS,
 BLISTER,
 BLOOD,
 BLOWPIPE,

CACHEXIE,
 CAJEPUT OIL,
 CALCULUS,
 CALLUS,

Page.

BLUNT HOOK,
 BONES,
 BORAX,
 BOUGIE,
 BRAIN,
 BRANDY,
 BREAD,
 BREAST,
 BRONCHITIS,
 BRONCHOCELE,
 BRONCHOTOMY,
 BUBO,
 BUBONOCELE,
 BURDOCK,
 BURNS,
 BURSÆMUCOSÆ,

CALOMEL,
 CALUMBA,
 CAMOMILE,
 CAMPHOR,

Page.		Page.
CANCER,	CINCHONA,	
CAPILLARIES,	CIRCULATION,	
CAPSICUM,	CIRCUMCISION,	
CARBON,	CLAP,	
CARBUNCLE,	CLAVUS,	
CARDIALGIA,	CLIMACTERIC,	
CARDITIS,	COAGULABLE LYMPH,	
CARIES,	COFFEE,	
CARMINATIVES,	COLCHICUM,	
CARTILAGE,	COLD,	
CASCARILLA,	COLIC,	
CASTOR,	COLLYRIA,	
CASTRATION,	COMBUSTION,	
CATALEPSY,	COMPRESSION,	
CATAMENIA,	CONCEPTION,	
CATARACT,	CONCUSSION,	
CATARRH,	CONFECTION,	
CATECHU,	CONGESTION,	
CATHARTICS,	CONIUM,	
CATHETER,	CONJUNCTIVA,	
CAUSTICS,	CONSTIPATION,	
CELLULAR MEMBRANE,	CONSUMPTION,	
CEPHALEA,	CONTAGION,	
CEPHALITIS,	CONTORTION,	
CERATE,	CONTRACTION,	
CEREBELLUM,	CONTUSION,	
CEREBRUM,	CONVULSION,	
CHANCER,	COPAIBA,	
CHARCOAL,	COPPER,	
CHEMOSIS,	CORNEA,	
CHERRY,	CORNS,	
CHEST,	CORPULENCY,	
CHICKEN POX,	CORYZA,	
CHILBLAIN,	COUCHING,	
CHILDBED,	COUGH,	
CHLORINE,	COSTIVENESS,	
CHLOROSIS,	COWHAGE,	
CHOCOLATE,	COWPOX,	
CHOLERA,	CRAMP,	
CHORDEE,	CRANIUM,	
CHOREA,	CRAZINESS,	
CHYLE,	CREPITUS,	
CHYME,	CRISIS,	
CICATRIZATION,	CROSS BIRTH,	
CICUTA,	CROTCHET,	

CURE,
CUTANEOUS DISEASES,
CYNANCHE,

DIAPHRAGM,
DIARRHCEA,
DIATHESIS,
DIET,
DIGESTION,
DIGITALIS,
DISCUTIENT,
DISLOCATION,
DISSECTION,
DISTORTION,
DIURETIC,
DIZZINESS,
DRESSING,
DROPSY,
DROWNING,

	Page.
DUMBNESS,	EXTINCT
DURA MATER,	EXTIRPATION
DYSENTERY,	EXTREMITY
DYSMENORRHOEA,	EYE
DYSPEPSIA,	HYALID

	Page.
DYSPHAGIA,	EXTIRPATION
DYSPHONIA,	EXTIRPATION
DYSPNOEA,	EXTIRPATION
DYSURIA,	EXTIRPATION

EAR,	FORGETS
ECHYMOSIS,	FRACURE
ECSTASY,	FRIGID
ECTHYMA,	FRONT
ECTROPIUM,	FRONT
ECZEMA,	FRONT
EFFLORESCENCE,	FRONT
EFFLUVIUM,	FRONT
EFFUSION,	FRONT
ELATERIUM,	FRONT
ELECTRICITY,	FRONT
ELECTUARY,	FRONT
ELEPHANTIASIS,	FRONT
EMACIATION,	FRONT
EMBROCATION,	FRONT
EMBRYO,	FRONT
EMETIC,	FRONT
EMMENAGOGUE,	FRONT
EMOLLIENT,	FRONT
EMPROSTHOTONOS,	FRONT
EMPYEMA,	FRONT

EMULSION,	FRONT
ENCYSTED TUMORS,	FRONT
ENDEMIC,	FRONT
ENEMA,	FRONT
ENTERITIS,	FRONT
ENURESIS,	FRONT
EPIDEMIC,	FRONT
EPILEPSY,	FRONT
EPILOCELE,	FRONT
EPISTAXIS,	FRONT
EPSOM SALTS,	FRONT
ERGOT,	FRONT
ERUCTION,	FRONT
ERYSIPELAS,	FRONT
ERYTHEMA,	FRONT
ESCAROTIC,	FRONT
EVAPORATION,	FRONT
EXCITEMENT,	FRONT
EXCITABILITY,	FRONT
EXCRESCENCE,	FRONT

	Page.
EXFOLIATION,	418
EXCRETION,	419
EXOSTOSIS,	420
EXPECTORANT,	421
EXPECTORATION,	422

FAINTING,	423
FANATICISM,	424
FASTING,	425
FAT,	426
FATUITY,	427
FEBRIFUGE,	428
FERMENTATION,	429
FEVER,	430
FIBRE,	431
FIBRIN,	432
FIDGETS,	433
FISTULA,	434
FLATULENCY,	435
FLEXIBILITY,	436

GALBANUM,	437
GALL-BLADDER,	438
GALL-STONE,	439
GANGLION,	440
GANGRENE,	441

	Page.
EXTRACT,	442
EXTRAVASATION,	443
EXTREMITIES,	444
EYE,	445
EYELID,	446

FLOODING,	447
FLUID,	448
FLUTTERING,	449
FŒTUS,	450
FOLLY,	451
FOMENTATION,	452
FOOD,	453
FORCEPS,	454
FRACTURE,	455
FRICTION,	456
FUNCTION,	457
FUNDAMENT,	458
FUNGUS,	459
FURUNCLE,	460

GARGLE,	461
GASTRIC JUICE,	462
GASTRITIS,	463
GELATINE,	464
GENERATION,	465

	Page.
GENTIAN,	127
GESTATION,	128
GLAND,	129
GLOTTIS,	130
GLUTTONY,	131
GOITRE,	132
GONORRHOEA,	133
GOUT,	134

HÆMATEMESIS,	135
HÆMATOCELE,	136
HÆMATURIA,	137
HÆMOPTYSIS,	138
HÆMORRHAGE,	139
HÆMORRHOIDS,	140
HANGING,	141
HARELIP,	142
HEAD,	143
HEADACHE,	144
HEART,	145
HEARTBURN,	146
HEAT,	147
HECTIC,	148
HEMICRANIA,	149
HEMIPLEGIA,	150
HEPATITIS,	151
HERNIA,	152

ICE,	153
ICELAND MOSS,	154
IDIOSYNCRACY,	155
IDIOTISM,	156

	Page.
GRANULATION,	157
GUIACUM,	158
GUM,	159
GUMBOIL,	160
GUMS,	161
GUN SHOT,	162
GYMNASTIC,	163

HERPES,	164
HICCOUGH,	165
HIP,	166
HIRUDO,	167
HOOPING COUGH,	168
HOOR-GLASS CONTRACTION,	169
HUNGER,	170
HYDATID,	171
HYDRAGOGUE,	172
HYDROCELE,	173
HYDROCEPHALUS,	174
HYDROMETRA,	175
HYDROTHORAX,	176
HYMEN,	177
HYOSCIAMUS,	178
HYPOCHONDRISM,	179
HYPERTROPHY,	180
HYSTERIC,	181

ILEUS,	182
IMBECILITY,	183
IMPETIGO,	184
INCONTINENCE OF URINE,	185

	Page.
INCUBUS,	
INDIGESTION,	
INFLAMMATION,	
INFLUENCE,	
INFUSION,	
INJECTION,	
INOCULATION,	
INSANITY,	
INSPIRATION,	
INSTINCT,	
INSTRUMENTS,	
INTERMENT,	

JALAP,
 JAUNDICE,
 JAMES' POWDER,
 JOINT,

KIDNEY,
 KINO,
 KNEE,
 KNIFE,

	Page.
INTERMITTENT,	
INTESTINE,	
INTUSSUSCEPTION,	
INVERMINATION,	
INVERSION,	
IODINE,	
IRIS,	
IRON,	
IRRITABILITY,	
ISCHURIA,	
ISSUE,	

LABOR,
 LACERATION,
 LACTEAL,
 LACTUCA,
 LARYNX,
 LEAD,
 LEECH,
 LEG,
 LEPROSY,
 LETHARGY,
 LEUCOMA,
 LEUCORRHOEA,
 LICHEN,
 LIFE,
 LIGATURE,
 LIGHTNING,
 LINIMENT,
 LIPPITUDO,
 LIQUID,
 LIQUOR,
 LISPING,
 LITHONTRIPTIC,
 LITHOTOMY,

[illegible]

MUCUS,
 MUMPS,
 MURIATIC ACID,
 NARCOTIC,
 NAUSEA,
 NECK,
 NECROSIS,
 NEGRO,
 NEPHRITIS,
 NERVE,
 NEURALGIA,
 NEUTRAL SALTS,
 NITRE,
 ODONTALGIA,
 OESOPHAGUS,
 OIL,
 OMENTUM,

[illegible]

Page.	
Page.	
15	OSTIFICATION,
15	OSTEO SARCOMA,
15	OTITIS,

PAIN,	
PALATE,	
PALPITATION,	
PALSY,	
PANACEA,	
PANCREAS,	
PAR VAGUM,	
PARACENTESIS,	
PARAPHIMOSIS,	
PARAPLEGIA,	
PARONYCHIA,	
PAROTITIS,	
PARURIA,	
PASSION,	
PATHOLOGY,	
PEARL ASH,	
PECTORILOQUY,	
PEDILUVIUM,	
PELVIS,	
PENIS,	
PERCUSSION,	
PERINEUM,	
PERIOSTEUM,	
PERIPNEUMONY,	
PERITONEUM,	
PERITONITIS,	
PERSPIRATION,	
PERTUSSIS,	
PHARMACY,	
PHIMOSIS,	
PHLEBITIS,	

Page.	
Page.	
15	OVARIA,
15	OXIDE,

PHLEBOTOMY,	
PHLEGMASIÆ,	
PHLEGMON,	
PHLEGMATIA DOLENS,	
PHTHISIS,	
PHYSIOLOGY,	
PILE,	
PILL,	
PIMPLE,	
PLACENTA,	
PLAGUE,	
PLETHORA,	
PLEURITIS,	
PLEURODYNE,	
PNEUMONIA,	
PNEUMOTHORAX,	
PODAGRA,	
POISON,	
POLYPUS,	
PORRIGO,	
POTASH,	
PREGNANCY,	
PRESENTATION,	
PROCIDENTIA,	
PROSTATE,	
PRURIGO,	
PRURITUS,	
PRUSSIC ACID,	
PSOAS ABSCESS,	
PSORA,	
PSORIASIS,	

Page.		Page.	
PSOROPHTHALMIA,	OVARIA	PUPIL,	OSTEOCLASTS
PTERYGIUM,	OXIDE	PURPURA,	OSTEOCLASTS
PTYALISM,		PUS,	OTITIS
PUERPERAL FEVER,		PUTREFACTION,	
PULSE,			
	PHLEBOTOMY		PAIN
	PHLEGMASIS		PALATE
	PHLEGMON		PALPITATION
	PHLEGMATIA DOLENS		PALSY
	PHTHIRIS		PANACEA
	PHYSIOLOGY		PANCREAS
	PILE		PAP VAGUM
	PILL		PARACENTESIS
	PIMPLE		PARAPHIMOSIS
QUARTAN,	PLACENTA	QUININE,	PARAPLEGIA
QUASSIA,	PLAQUE	QUINSEY,	PARONYCHIA
QUERCUS,	PLETHORA		PAROTITIS
	PLEURITIS		PARURIAS
	PLEURODYNE		PASSION
	PNEUMONIA		PATHOLOGY
	PNEUMOTHORAX		PEARL ASH
	PODAGRA		PECTORIL OGUT
	POISON		PELVIVUM
	POLYPUS		PELVIS
	PORRIGO		PENIS
	POTASH		PERCUSSION
	PREGNANCY		PENNIUM
RABIES,	PRESENTATION	RENAL,	PERIOSTITIS
RACHITIS,	PROCTITIS	RESIN,	PERIPISTIMUS
RANULA,	PROSTATE	RESOLUTION,	PERITONITIS
RASH,	PRUNICE	RESPIRATION,	PERITONITIS
RATTLE,	PRUNITIS	RESTLESSNESS,	PERITONITIS
RECTUM,	PRUNICACID	RESUSCITATION,	PERITONITIS
REGIMEN,	PSORA ABSCISSA	RETENTION,	PHARYNX
RELAXATION,	PSORA	RETINA,	PHARYNX
REMITTENT,	PSORIASIS	RHEUMATISM,	PHLEBITIS

	Page.
STERTOR,	ROSTOL
STETHOSCOPE,	RUBCOL
STOMACH,	
STONE,	
STRABISMUS,	
STRAMONIUM,	
STRANGURY,	
STRICTURE,	
STROPHULUS,	
STUTTERING,	
STYPTIC,	
SUBSULTUS,	
SUDORIFIC,	
SUFFOCATION,	
SUGAR,	
SULPHUR,	BRIN
	SKUL
	SLEED
	SMALL FOX
	SMELL
	SNEEZING
	SNUTE
	SODA
	SOLD
	SOLUTION
	SOLVENT
	SOUND
	STACELUS
	STAIN
	SPEECH
	SPHINCTER
	SPIGELIA
	SPINA RUPEA
	SPINAL MARKOW
	SPINE
	SPRINT
	SPRINK
	SPRINT
	SCULLS
	ST. ANTHONY'S FIRE
	STAPHYLIONA
TABES,	STAY
TÆNIA,	STAY
TARAXACUM,	STAY
TASTE,	STAY
TEETH,	STAY

	Page.
SULPHURIC ACID,	118
SURGERY,	119
SUSPENDED ANIMATION,	120
SUTURE,	121
SYMPATHY,	122
SYNCOPE,	123
SYNOCHA,	124
SYNOCHUS,	125
SYNOVIA,	126
SYPHILIS,	127
SYRINGE,	128
SYRUP,	129
SWEAT,	130
SWIMMING,	131
SWOON,	132
	133
	134
	135
	136
	137
	138
	139
	140
	141
	142
	143
	144
	145
	146
	147
	148
	149
	150
	151
	152
	153
	154
	155
	156
	157
	158
	159
	160
	161
	162
	163
	164
	165
	166
	167
	168
	169
	170
	171
	172
	173
	174
	175
	176
	177
	178
	179
	180
	181
	182
	183
	184
	185
	186
	187
	188
	189
	190
	191
	192
	193
	194
	195
	196
	197
	198
	199
	200
TEETHING,	201
TEMPERATURE,	202
TENDERNESS,	203
TENDON,	204
TENESMUS,	205

Page.	
TESTICLE,	VENIS.
TETANUS,	VENTRIQULE.
TETTER,	VERMIN.
THERAPEUTIC,	VOICE.
THERMOMETER,	WOMAN.
THIRST,	WOMAN.
THORAX,	
THROBBING,	
THROMBUS,	
TIC DOLOREUX,	
TIN,	
TINCTURE,	
TINEA,	
TISSUE,	
TONGUE,	
TONIC,	

WHITOW,	
WIND,	
WOMB,	
WORMS,	
WOUND,	
WRYNECK,	

ULCER,	WINE.
ULCERATION,	WINE.
UMBILICUS,	WINE.
UNGUENT,	
URETHRA,	

Page.	
TONSILS,	VACCINATION.
TORPOR,	VAGINA.
TOUCH,	VAGUE.
TOURNIQUET,	VAGUE.
TRACHEA,	VAGUE.
TREMOR,	VAGUE.
TREPANNING,	VAGUE.
TRICHIASIS,	
TRISMUS,	
TROCHES,	
TRUSS,	
TUBERCLE,	
TUMOR,	
TWINS,	
TYMPANITES,	
TYPHUS,	

WARMTH,	
WART,	
WATER,	
WEA,	
WHITES,	
WHITE SWELLING,	

URINARY CALCULUS,	WINE.
URINE,	WINE.
URTICARIA,	WINE.
UTERINE HÆMORRHAGE,	
UTERUS,	

Page.		
	VACCINATION,	TONSILS
	VAGINA,	TORTOR
	VAPOUR,	TOUCH
	VARICELLA,	TOURNIGURT
	VARICOCELE,	TRACHEA
	VARICOSE,	THORAX
	VARIOLOID,	TREANNING
		TRICHIASIS
		TRISMUS
		TROCHES
		TRUSS
		TUBERCLE
		TUMOR
		TWINS
		TYMPANITES
		TYPHUS

WAKEFULNESS,
 WART,
 WATER,
 WEN,
 WHITES,
 WHITE SWELLING,

YAWS,
 YELLOW FEVER,
 YELLOW GUM,

Page.		
	VEINS,	TESTICLE
	VENTRILOQUISM,	TETANUS
	VERMIFUGE,	TETTER
	VOICE,	THEATRUM
	VOMICA,	THROMBOSIS
	VOMITING,	THYROID
		THORAX
		THROBBING
		THROMBUS
		THE DOLOREUX
		TIN
		TINCTURE
		TINEA
		TISSUE
		TONGUE
		TONIC

WHITLOW,
 WIND,
 WOMB,
 WORMS,
 WOUND,
 WRYNECK,

ZINC,
 ZINGIBUR,
 ZONA,

ULCER
 ULCERATION
 UMBILICUS
 UNGUENT
 URETHRA

Notes taken from Dr. Colhoun's Lectures

Material Medica 1835

November

Modus Operandi of Medicines

Some medicines act generally on whole system by being applied to any part of body as *Stimulants*
 Others act locally by being applied to a particular part
 Are medicines absorbed some indirectly are
 Lymphatic & venous systems are the systems which absorb
 Medicines also act by imbibition or soaking
 & they also sometimes act by sympathy

Blood Letting

There are 7 modifications of pulse 1st A hard pulse 2nd full
 3rd sudden or abrupt 4th quick 5th rapid 6 variable
 7. intermittent pulse

Hardness indicates great force of heart's action

Cases which contra indicate blood letting

A pale & sallow countenance 2nd pain in breast with a sensation
 of stricture not attended with cough when patient sits up
 3rd where the bowels are loaded with feculent matter keeping up
 the irritated excitement
 4th where there is pain in head & this increased when patient
 sits up

Convulsions sometimes occur after fainting from bleeding

Children do not bear large blood lettings without the exists local inflammation
 in some important organ but they bear it frequently

Cathartics

December 1st 1835

Medicines that have the effect of producing evacuation from bowels when
 exhibited in sufficient doses Cold suddenly applied produces
 evacuations also does certain articles of diet

It becomes necessary to evacuate the bowels in every violent disease

Chronic constipation may sometimes be relieved by rising early
 taking exercise in proportion to strength diluents also assist

Acid watery bitter sweet & fatty substances have a tendency to produce evacuation
 walking on cold ground also has a fine effect

Cathartics

Mercury in chronic constipation has a fine effect not only by producing
alterative effects but also by producing pancreatic & biliary secretion.
Cathartics are of 2 kinds viz. Drastic & mild

Notes taken from Dr Colhoun's 1836 Lectures on Materia Medica

Nov 15th Discovered by Dysc in 1645
Cephaelis Ipecac. v. America
Root is annular in its cortical part This is its
distinguishing character
Emetic has been found in the largest quant in brown variety
There is however only 2 per cent diff
Its virtues ^{depend} on brilling It should not be
exposed to air or light (old)
Process for prep. Emetic or any other alk.
You add ether to remove fatty matter
Alkaloids are gen insoluble in cold water, little sol.
in warm & very soluble in hot alcohol
Precipitation of gall antidote to ipecac.
Ipecac Dose in powder 15 gr to $\frac{1}{3}$ as an emetic
Dose in val. prep. Vinum Ipecac. squibb.
vinegar oramel &c are given in dose of oz as
an emetic $\frac{1}{3}$ is a diaphoretic dose
Ipecac is safe and efficacious vomit
gum arabic takes away its nauseous taste
Qid in small doses it has a fine effect in allaying
the appetite
It is better than tartar emetic in debilitated cases
or in an irritable state of stomach and bowels
It is suited partic. to chronic dysentery
Boil it with ether afterwards filter it

Nov 17th *Specac.* - Emetic Antidotes infusions of
galls Acetic acid

Specac sometimes combined with bitter extracts as gentian
It is found to suspend its emetic effects

Hemorrhage It is sometimes given in these cases

Dyspepsia very val. in slight cases by increasing tone of stomach
& keeping bowels open It is in many cases owing to costiveness

Sulphur Livi - White vitriol
used where large doses of opium has been swallowed
I throw $\frac{1}{2}$ $\frac{3}{4}$ in tumbler of water & administer it
Ashes administered decompose it & may be used as an antidote
Carbonates of alkalis are antidotes to it

Dose 10 gr to $\frac{1}{2}$ as an emetic. It is mild in its operation
a trill

Ophthalmia val in these cases after inflamed symptoms have been
Howallus It is very valuable in some cases

Tartar Emetic

It is decomposed very soon in form of solution = 24h

Nov. 18. made by boiling protoxide of antimony with cream of tartar
Incompatibles Decoc of bitter & astringent vegetables
one of best Tar. E - 1 gr
purgative & Soda or Mag. 3
is the fol. Ag. font. $\frac{1}{2}$ pt

Dose wine glass full every hour or 2
As nauseant med. $\frac{1}{4}$ gr

$\frac{1}{2}$ gr every 15 m. in tumbler of water
sometimes it is given in form of powder = objection
Tolerance of medicine increased by give it with opium
I would not go beyond a gr at a dose
This medicine may be give in quant of 23 injected in
to rectum

Effects vomiting purging increase of urine
& sweat

Emetics

Nov. 18th 1836

Tartar Emetic Wine 2 grs to the oz is the proportion used.
 Medium dose is 13

Medicinal
uses

Antimony sometimes salivates

It is given in fevers & in almost all inflam. diseases
 Interesting fact - It cures inflam. of veins when given
 in nauseating doses

Nicotiana Glauca

Nov. 21st

Vomiting is its most general effect. It prostrates
 the system. Best applied to surface in form of

Colic poultice. It relieves colic in a very happy manner.
Tetanus has been cured by it

A. Cholera Dr G. M. Clellan cured a case by this remedy
 In debilitated persons it must be given in small doses

Dose 15 to pint water

Dysentery Applied externally in form of decoction

Croup do do to throat

It also produces abortion in some animals

Lobelia Inflata

Its virtues reside in almost all parts of plant

It produces nausea vomiting great debility &c

It is one of our most valuable emetics

Asthma most valuable remedy. It is caused by any
 thing that obstructs the lungs

Yable spoonful of tincture at a dose has relieved it

It should be given in August. It may be given in
 infusion or tincture

Equal parts of Lobelia tincture & syrup of squills
 cures acute bronchitis in the most happy manner

Blue Vitrol - Bi sulphate of Cop

Dose 5 to 15 grs. It is prompt & does not produce

much debility

Antidote Galls is the best antidote

Salts of copper are all poisonous

Emetics

Nov. 22nd 1836

Mustard give in dose of a teaspoonful largely diluted with water has the effect of producing vomiting
 Horseradish, Inner bark of elder &c are good emetics
 Emetics are instrumental chiefly in lessening the action of system
 & have 5 diff effects upon the system

Nov. 28th **Cathartics** Nov 28th 1835
 Drastic Elettarium, Eup

Active Jalap. Senne

Laxative sulphur

1st of Drastic Cathartics
 Elettarium. It is contained in pods around seeds Insoluble in cold or warm water Dose of Clutternucks prep 1/8 to 1/10 gr Powder of leaves

Elettarium Dose 1/2 to 1/6 gr

Recommended in dropsy In large doses it produces pain inflammation Spasms Coma Convulsions

Rheumatism of Acute character = Recommended

Croton Tiglium

oil of seeds is the part used It has a pale redish brown colour It contains 15 parts of an acrid principle which is soluble in alcohol

Dose 1 gr.

Medicinal you can use

It is one of the best rubefacients that

uses

gum Arabic

It may be give in mucilage of

Dropsy become in Croton Tiglium rubbed up with gum arabic and applied to travel in quant of 6 drops

In

Euphorbia Lathoris Mole plant

It has all the active prop of croton oil give rubbed up with gum Arabic & sugar
 Sarsaparilla is an active purgative

Nov. 28th — *Cucumis Colocynthis*

— *Colocynthis* Active principle
 Medicinal) Palsy of extremities & bladder
 uses } official preparations Comp. Ex. *Colocynthis* which
 may be given with calomel Dose 5 gr. to $\frac{1}{3}$

— *Black Hellebore*
 It is a hydragogue drastic cathartic Not of much use

— *Aloes*
Socotorine - *Hepatic* *Cape aloes*
 It softens under the fingers
Cape Aloes commonly met with in market & is very good
 It is the only one which we traced strictly
 Nearly insoluble in water Alcohol is a good solvent
 Medicinal) It operates strongly on rectum
 uses } Menorrhoea Habitual costiveness
 Dose Medium dose 10 gr — It is one of the best purgatives
 in cases of debility attended with enlargement of abdomen
 Fructus dose $\frac{1}{2}$ to $\frac{1}{2}$ 3

December 1st

— *Scammony*
 Exceedingly drastic cathartic *Allypo scammony* best
 Dose 3 to 15 gr

— *Gamboge* obtained from *Stigmatis gam.*
 It is a drastic cathartic Combined with *Cremor Tartar*
 it has a good effect as purgative
 Compound pills only official prep used

— *May-Apple* *Todophyllum* *Littatum*
 Given in same dose as *Galap*
 Perennial plants should be gathered in winter before
 they flower. Biennial plants during 2nd year
 while in flower
 all the above articles are drastic in their effects
 They should not be given when there is pain
 or acid in stomach or bowels

Purgatives December 10th 1836

Active Cathartics or middle variety Jalap.

Alcohol is prop. solvent May be combined with Calomel or cream tartar

Medicinal uses Combined with cream tartar in teaspoonful doses is one of best purgatives in dropsy & disease of joint

Resin of Jalap - adulterated by guaiacum

Rhubarb

Medicinal uses To be good it should be heavy & solid Water is the solvent Combined with aloe it forms best purgative in Dyspepsia

To relieve acidity it should be combined with magnesia

Combined with aromatics valuable in gout

It tinges the urine of dark brown colour

Dose of 40 gr is less apt to gripe than 20 gr

Aromatic syrup of Rhubarb is a good prep

Senna

It is adulterated with box leaves which may be known by applying heat - it causes the 2 plates of box leaf to separate

It is also adulterated with a heart shaped leaf

It is one of the most common purgatives in children

Cassia Merilandica

It is a purgative of great power and used in same way & in same dose as Senna

Dec. Ind.

White Balsam - Bull's head - 10 grs - good to oblige

Caster oil - castor oil with mineral oil - 10 grs - good to oblige

oil, clove oil - known by longish leaves in strong alcohol

It has been given in colic & Diets - 10 grs

Calomel - 10 grs - 10 grs of mercury

Dose 8 grs - 1/2 grs - 1/4 grs - 3 to 4 better effect

opening purgative in the first young persons

Purgatives

Calomel - sometimes adulterated with corrosive sublimate
 removed by agitating it with water & to the water add lime water
 & the Calomel Medicament

They cause some stools. Combined with tartar emetic
 they are one of our most useful remedies in effluvia poisons

Glauber's salt

Dose $\frac{1}{3}$ to $\frac{2}{3}$

Epispurus salt resembles E. C. C.
 acid white vitriol. Aequal parts of Sulfate of Mag. &
 common salt form

Sal solution of Mag. 7.3. 2 Mag. bitter & E. C. C. Acid

Powder or tartar powder is a violent effect

Phosphoric acid. Dose 1.3

Seedling powder

Sulphur obtained from Italy & Rome
 to be kept in a temperature to detect arsenic in it
 Sulphur is the best. Dose 1.3 to 2.3

Dec 5th — Cassia Fistula

In dose of an oz or more is an excellent purgative

Tamarinds

Dose same as last article

Peach

Strong infusion of peach
 2.3 to first. It produces an emetic & purgative of
 Olive Oil

Adulteration. It is obtained by ~~appreciation~~

known by coagulating at temperature of
 Adulterated with linseed or poppy oil. Dose 2.3

In poisoning from reptile you must treat the case
 scientifically. Examine the pulse. If the
 pulse is scarcely to be felt alcohol is the proper
 remedy. If you rely upon the specifics that
 have been recommended your patient will surely
 die

Magnesia Dec - 5th 1836

It suspends camphor & opium in water
Carbonate of magnesia is antidote to sulphuric acid
Pure magnesia is not prop to be giv in sick case as it generate
intense heat in uniting with S. Acid
Carbonate may be tested by adding vinegar when considerable
effervescence will ensue

Dec 6 Sulph. A. has been known to kill in dose of 3
Oxalic acid known by sour taste & by crystals res of Magnesia
& by its taking away the stain of ink
Magnesia dissolved in carbonated water
It has been giv in various ways may be giv with eg.
juice of orange peel & giv in dose of table spoonful
Magnesia triturate
Adulterations chalk

Manna
Giv to children in dose of from 3 to ʒ 3

Blue Pill

They operate at a period of 10 or 12 hours

White & Black Mustard Dose is
a table spoonful ev morning taken whole
or a little water

Oil of butter is a valuable purgative

Diaphoretics

Dec 23rd 1836

Mitre Antimony Species.

Camphor may be giv with antimony Dose ʒ 5. or ʒ 9

It may be taken with warm vinegar

Combined with opium valuable in typhoid

Common vinegar is valuable as a phlegma point of view
combined with other medicine in treat of febrile diseases

Adulteration with sulphuric acid known by dipping Cornish
Rage into it when it becomes affixed

Vinegar when giv alone when giv warm is a valuable adjuvant
but its combination more valuable

Citric Acid Dec 23rd 1836

when taken alone is debilitating with the salts

it is one of the best debilitating diaphoretics

20 grs of Potash salt by 15 gr Citric acid

— ammonia — by Citric acid &c

Acqui carbonate of ammonia

10 grs with 2 grs of ammonia given at night

Eupatorium Perfoliatum has the same operation

bitters & acts as a diaphoretic

Give in strong

tea $\frac{1}{2}$ $\frac{3}{4}$ to 1 gr water

Andalapsus Sabiorea

Cucurbit Laseparilla

Dose 10 gr

Pruriger venous & sometimes salivator

It may be given in cases in which system is not very

Daphne Mezereum

exter applied it blisters the skin

23 bark of root in 3 pts of water down to 2

Smilax Laseparilla

It produces an acid upon the tongue which is a test of its goodness

As a mild diaphoretic it will do good in all cases

where a diaphoretic is required Dose 10 grs

Dec 25th Soda powders 25 Tartaric Acid & 10 grs of bi-carbonate of soda put up in separate papers = They furnish a good substitute for soda water

Tartaric Acid & a constit of seedling powders = 23 of tartarate of potash & soda (Rochelle salt) & 23 of bi-carbonate of soda put up in a white paper & 35 gr Tartaric acid contained in a blue one

Expectorants

no specific in the class of expectorants

Dec 30th 1836

Diuretics

Jan 5th 1837

Digitalis given dose of 10 or 15 grs may for 3

Veratrica

Medicinal uses

Erosy 104 gr

to 3 of lard & rubbed upon skin

Dose $\frac{1}{2}$ gr

Pindar Jan 25/1826

Because I suppose by G. Byron to be
 a set off and means against the par
 ticularist points in woman
 I might this be true you will excuse it for
 I am very young & as smart as the rest of you
 I know it
 the 3rd
 line by
 Byron
 I mean
 by me
 M. D. G.

Why is backhear cake like a Caterpillar
 Why is the letter S in the word Citizen like a persons nose
 There is a thing in Amsterdam in Rome it doth appear
 Its twice in every moment not once in seven years
 Christ never saw Washington seldom saw what we
 see every day

Why are Ladies like Coach Drivers? (males)
 Because their first object is to secure the males
 Why by the law of retaliation have you a right to pick a
 painter's pocket

I am a word of one syllable short & plain, which spells
 backwards & forwards the same it expresses the warmest
 sentiments fresh from the heart & to beauty lays principle
 claims

Why is a room full of married people like an empty room

Jan 5. Neutral Diuretics: those which neither increase ^{the action of heart & vessels} or decrease

Carbonates of Potash & Soda
 Spiritus Ether Nitricus Sweet Spirit of nitre
 Given in dose of 1, 2 or 3 tea spoonfuls
 Sulphuric & Vitric Acids
 Dose 10 drops 3 times a day taken large quant of water & taken
 through a quill

Murated tincture of Iodine Dose 10 or 15 drops
 10 minutes until it produces nausea
 Erigeron Heterophyllum Dose Gill of decoction

Iodine Though it has other properties
 deserves a place among the ~~diuretics~~ It is dissolved
 in 7,000 parts of water When boiled in a solution
 of potash it forms the hydriodate of Potash
 Ioduret of lead may be given in dose of 10 gr ^{to 1/2 gr sufficient}
 to begin with Tincture of Iodine 15 gtt which contains
 1 gr of iodine

best - its best given in form of Hy. of Potash with 1 gr of iodine
 dissolved in water & given in dose of 10 gr in ^{total 15 gr dry}
 Hydriodate of potash dissolves in the quant of water that is contained in it will
 keep in this way for 2 months & years

Iodine - Jan. 5th. 1837

Iodine should not be given in greater than $\frac{1}{2}$ gr in course of day
in chronic diseases

Bath

8 gr to 3 to oz laud is the proportion for outward

Exhibit in moderate doses it increases appetite & produces
diuresis

Medicinally Scrophulous is often relieved by it Goitre
Uses Drops 1 gr Iodine 8 gr. Hydriodate of Potash in day =
I gave $\frac{1}{2}$ gr of Hydriodate dissolved in water in a day

Diet Jan. 6th 1837

Man is omnivorous Water Exercise & diet
are three great Physicians

Soup is not so digestible as supposed

Fibrine - muscle is composed chiefly of fibrine. = $\frac{7}{11}$ water + fibrine

There is only $2\frac{1}{2}$ of soluble in water. Therefore

what is called essence of beef is not nourishing

Gelatinous a jelly - distinguishable by being precipitated by tannin

Albumen - specimen is the white of egg

Oil: composed of oleine & stearine

Vegetable Elements

Fecula

Diet in relation to its nutritive qualities

Animal food is the most nourishing

Fresh beef & mutton most digestible should be
broiled & underdone

Broth & all young meats is the least digestible

& at same time less nourishing

Those that most resemble meat are the

vegetables. Those which contain sugar

are more digestible than the watery

Fecula is the chief ingredient in the Tapioca & sago
Starch is the same as Fecula

Diet

Jan 9th 1837

Pea & bean are their nutritious qualities to March
 Beef Mutton most digestible & nourishing
 Fowl next Lamb veal &c
 Oysters a diet of oysters is debilitating
 Fish is less nourishing than Flesh
 Eggs & oysters are not more stimulating than vegetable food
 The diff kinds of fermented & spirituous liquors have the
 effect of assisting digestion
 Wines of time become acid to remove which lead is put
 into it Sulphuretted hydrogen is the test

Individual articles

Milk
 one of the most agreeable articles in certain diseases It contains but
 little nourishment Eg parts of lime water & milk
 suits well the stomach Whey is still less nourishing
 & is therefore a good remedy to deplete the system Influenza
 of stomach &c. S. J. milk 1824

Chronic disease can be cured by diet alone
 Jan 9th 1837 Rennet whey is the best diet in inflammatory diseases

Articles constituting low diet
 1st Ising-glass It is almost pure gelatine: 98 in 100
 It should be in the form of staple that obtained
 from the intestines is

Stecula = rice, arrow root, sago are composed chiefly
 rice 96 in 100

Chocolate adulterated with fat & sugar

Cinnamon salt

Pure Water contains nourishment

Nitrate silver Tartar emetic should be dissolved in pure water

Fresh rain water is quite pure & will answer

all the purposes in place of distilled water

Honey is sometimes poisonous

Sugar for the purposes of pharmacy answers every purpose
 with honey
 sometimes adulterated by flower Known by dissolving it
 in water

Diet Jan 10th 1837

Tea & coffee dispeptions should leave of both They use it too
Tea constipates & coffee does not
Tea is very low as nourishment but has considerable stimulant
and narcotic effects

Head ache comes on in consequence of leaving off tea & coffee
& is owing to low action

Muslinum one species is healthy & forms an agreeable ^{rich} morsel
rose red strated structure below

If they purgess they should be thrown off by a vomit & a purge &
if the pulse be weak, ether spiritum &c

Indian Corn

Diet as a preventive of
disease

What will prevent will cure is a true maxim

Animal diet will cure intermittent fever

mutton beef & fowls therefore form a good prevention

Bark will also prevent ague

Jan 10th 1837, Oat meal gruel is a very easy
of digestible

All kinds of fat long dried salted baked beef
are indigestible

Secula in all its forms has a tendency to ~~terrest~~ diarrhoea
Cold water in summer sometimes produces
pain Cramps Convulsions &c

Constipation relieved by diet & exercise
solid food lying in bed to long wearing flannel next skin
have a tendency to produce constipation

Those that ~~that~~ have a tendency to keep them open
Watery saccharine Acid bitter & oily

Brown bread have a fine effect in keeping bowels open
Milk & milk once or twice a day all have a tendency
to relieve it

Tomatoes change the stools black
Oysters & minceles where patient is unwell
very weak they should not be used

Notes Taken from Dr S Colman's Lectures on Diet Jan 12th 1837

Calumia Latifolia eaten by pleasant renders it poisonous
When taken into stomach reduces, in a few hours, the
pulse down to 40
Scurvy cured by meat of turtle which shows that a low diet
will cure this disease

Jan 12th 1837 Diet as a Restorative of Disease

Beef mutton Black meat food restore the strength
more rapidly than any other diet
Mutton is the most digestible of all kinds of food

Preparation of food

Roasting renders meats tender
Baking also confines the juices Boiling is the best
way of preparing it Slightly boiled meat is digestible
goul. ogt 1 to 3 water mix & boil it in pint water

Intacids Jan 13th 1837

Those who drink spirits are seldom subject to it if they
take plenty of exercise

In general all indigestible substances produce it
Vegetable food is liable to produce it Animal food is not
Muriatic is secreted during digestion & Acetic acid in small
quantities

Some potash soda

Carbonate of Potash or em. potash By passing
through lime it is freed of its acid

Bicarbonate of Potash bicarbonate

& bicarbonate are used to correct acidity

Carbonates of potash & soda unite with purgatives

have a good effect in correcting the stools

Sometimes stimulating gums or ammonia is used with

these carbonates

Ammonia aqua 20 or 30 grs

36
Lime Jan. 13th 1837

Used in form of lime water or carbonate of lime - Com. chalk
Chalk 13 hand. 123 9 Arabic 13 water 83
Charcoal is a good collector of acidity
Mercury enters up with magnesia is good Antacid

Astringents Jan 16th 1837

Vegetable Astringents Tannic acid is the astringent principle
It unites with oxide of antimony of tartar emetic & is therefore its antidote

Vinegar is no antidote as formerly supposed

Medicinal } Tannic acid is to astringents what gumme is to Cinchona
Uses } Has the effect of arresting hemorrhage
They act as blandants upon the gen system

Oaks - white oak black oak &c
Mineral Astringents

Acetate or Sugar of Lead
Weak solution 6gr to 3 water Strong 12gr to 3
Collyria used in strength of from 1 to 2gr
Dose 1gr with 5gr of opium

In large doses it produces inflame of stomach

Painters Painters will apply to you Wash hands
before meals change clothes freq. & when there is a tendency
to costiveness give castor oil

Antidotes Common Epsom salts is perhaps the best antidote
Jan. 19th Alum or Sulphat aluminous et potass.
10 gr every 2 hours It is by no means poisonous

In irritation of mucous membrane it is detrimous

23 in pt milk proportion for alum whey

Alum 5 gr
Sulphat 10 gr } 3 t. d.
2 or 3 gr in 1/3 oz collyria Ep bark 10 gr

Sulphur Zinc

Adulterated with iron copper & lead = Sulphuric Acid

Astringents Jan 19

Collyra = 1 or 2 grs to oz water

Sulphate of Copper

3 grs twice a day Sugars antidote to caps & files
 It is not an antidote Hemorrhage of Potts may be
 a valuable antidote Iron filings

Sulphuric Acid

It is one of the best rem in fever at that period when it is
 doubtful whether that fever requires stimulus or not
 Dose 10 grs 3 tds Antidote magnesia

Vegetable Astringents & vegetable bitters & tonics are
 as a general rule give in the dose of from 10 grs to 30
 Tonics Columbo

In cases of simple dyspepsia give a blue pill at night
 & in morning a wine glass full of infusion of Columbo Gentian
 Quassia &c Gentian

Phosphoric beakts Tremulous motion of hand blood
 shot ^{& staring} eyes Weak knees smacking of the lips as if the
 person was tasting something short dry cough; These moral
 habits are bad

Stimulants Jan. 20th.

48

45

46



Notes Taken from Dr. S. McClelland's Lectures
on
Obstetrics & Diseases of Women & Children

Notes taken from Dr. Dunglison's Lectures on Physiology

1836
Nov. 16th

Sympathies are
of three kinds 1st general 2nd continuous
3rd Contiguous Sympathy
Sympathies are transferred to membranes of analogous ^{structures}
Rheumatic inflammation of heart may exist without pain
Stomach sympathizes with uterus during pregnancy
If you apply child to breast soon after birth it brings
on after pains
Sympathies freq. become developed during disease
Sympathy of imitation
Abortion sometimes the consequence of labour observed in
another

Complexity of organization renders them the more
susceptible of infestation

Nov 17th

- 1 Glycistrophy or augmentation of function
- 2 Immulation
- 3 perboration of function
- 4 Suspension of the action for a time
- 5 Arritation

Diseases of stomach is very frequent

Disease of liver is very frequently secondary

Nov. 18th Stomach & heart more liable to disease than liver or spleen
Change in vital properties is for by change in
structure of the tissue or organs
In 177 cases that have the appearance of debility
are attended by increased action

Habit sensibility

Nov 21st The power of feeling & of moving responsive to such ^{feeling}
Functions - the action of an organ under the influence
of its vital properties

They are divided into Animal Natural & generation

Nervous System

Nov 22nd

Nervous system is divided into Cerebro-spinal axis & ganglionic systems of nerves

- | | | |
|--|--|---|
| 1 Regular | of all the spinal
&
one cerebral
viz | 5 th pair
suboccipital
7 cervical
12 dorsal
5 lumbar &
6 sacral |
| 2 Irregular
or
Superadded | 3 rd 4 th & 6 th to the eye
7 th to the face
9 th tongue
Glosso-pharyngeal to pharynx
Vagus to the larynx heart,
lungs & stomach
Phrenic to the diaphragm
Spinal Accessory to the muscles
of the shoulder
External respiratory to the
muscles of out sides of chest | |
| 1 Special sensibility | olfactory
optic
Lingual branch of 5 th pair
Auditory | |
| 2 General sensibility | 5 th pair
Spinal nerves
(posterior roots) | |
| 3 Voluntary motors | Spinal nerves (anterior roots)
Common oculo-muscular
External oculo-muscular
Hypoglossal
Pathetic, facial, glosso-pharyngeal
al. phrenico-gastric &
spinal accessory | |
| 4 Instinctive motors
or
respiratory nerves
of Sir C. Bell | | |
| 5 of vital association
& nutrition | Filicolumns & plexuses of the
Ganglionic System | |

69
 Nov 25th 1836

First link in the chain of fever is derangement in functions of organic nervous system
 2nd. Alterations in secretions 3rd derangement in sanguiferous system

Nov. 29th

Respiration

There is a kind of respiration going on by skin
 Body contains 32,000th part of pure air

Dec 7th Syrups have the effect of purifying the blood
 Hematosis = blood making

I believe that atmospheric air enters into circulation
 Carbonic acid is formed in the blood

Quant. of air consumed & quant of carbonic acid given off
 Dec 8th Can you decide in a court of justice independent of circumstantial evidence that the individual was either hanged or drowned by mechanical impediment or by irrepressible gases. You cannot distinguish one of these modes from the other

Insufflation is the best means to restore an asphyxiated person
 No water enters into stomach except while individual is drowning or after putrefaction has taken place

Circulation &c

In case of hemorrhage from translocation you should not allow the patient to drink freely of very thin fluids keep patient of light food & allow him acids that will increase the blood
 Sulfuric Acid S.G. blood 1080
 Increased the sp grav. is sometimes 1022

Dec 12th

Polyemia or Plethora

Anemia less quant of blood than natural
 we mean a smaller quant of blood as well as poorer that is it will be poor in red particles
 Brought on sometimes by excessive blood letting
 Serum consists of albumen with salts
 S.G. 1027 in health

70
Circulation Decr. 12th 1836

When pericardium is inflamed it produces a sound like cracking of leather
Healthy quant of liquid in pericardium There may be
2 oz without disease Usual quant tea spoonfull

Other

Notes taken from Dr. Revere's Lectures on Practice of Medicine

Nov. 5th Theory & Practice of Medicine divided into
Physiology, Pathology, Therapeutics

Physiology that branch of science which treats of the functions of the living animal body & of the powers by which these functions are exercised

Pathology the doctrine of diseases

Therapeutics that branch of medicine which treats of the operation of the different means for curing diseases & of the application of these means

Temperaments or more properly Constitutions

Forms of constitutions they are the Robust flaccid & ^{Scrofulous}

Robust persons - all who habitually enjoy good health & whose muscles feel firm under pressure

2nd class Their configuration is such as to indicate a delicate habit the muscles when grasped feel soft

3rd class Their complexions are usually fair; hair ^{gen} light eyes also ^{gen} light upper lip is protuberant body is of a delicate & curved

Scrofulous persons are not necessarily diseased

constitutions are sometimes hereditary

Inhabitants who live in cold & damp situations are especially prone to scrofulous constitutional affections

2 methods of treating diseases viz Expectant & Active

Expectant mode consists merely in the removal of causes which produced the disease

Active treat consists in the administration of medicines &c

Idiosyncrasy A peculiarity of constitution observed in a few individuals from particular agents

Diathesis Any particular state of the body
Classification of the organs & functions

They are arranged into Animal functions which consist of sensation thought & voluntary motion Vital - these are Digestion, ^{Respiration} Circulation & nutrition Natural & the Generative functions

Novemberth Richats Classification of the functions is into the
 Nutritive functions - digestion respiration circulation &
 nutrition 2nd the functions of relation 3rd the
 generative functions

Excitability - example When a child is born & respiration does not take
 place, by applying a little cold water or brandy ^{to the stomach} may ^{cause}
 the ~~child~~ ^{lungs} to expand This excitement
 is owing to excitability of system

Excitement the effects of the stimuli acting upon the excitability
 Every organ & texture has its peculiar excitability & its appropriate stimulus
 Diagnosis the science which delivers the signs by which a disease may be
 distinguished from another disease

Prognosis the foretelling the events of diseases from particular symptoms

The Distinctive Characters of Disease

Inflammation There are but 4 textures which do not inflame viz hair
 nails epidermis & enamel

Definition a local disease in which there is an accumulation of blood
 in part with a deranged state not only capillaries but also of
 all the other textures which go to make up the part & is attended
 with pain redness increased heat redness & tumefaction

Mode of distinguishing inflammation in internal parts
 When we have superadded to pain a morbid sensibility in the
 part ^{some times} with a tumefaction & a derangement of functions of the system
 you may be assured of inflammation in part

There is generally a symptomatic fever

Causes of Inflammation Predisposing & Exciting
 most common causes of local inflammation
 changes of temperature heat when above 100° F
 cantharides turpentine gums Arsenic corrosive sublimate
 mechanical injuries

Terminations of inflammation 1 termination Resolution
 with or without changes of structure 2 adhesion 3 formation
 of false membrane 4 Effusion Suppuration 6 Ulceration
 7 Gangrene

Inflammation

77

1st of resolution without change of structure

Termination in Gangrene

Constitutional symptoms of mortification. A sudden alterations of pain in the part, features are very much contracted. There is great appearance of distress. Pulse becomes frequent & weak subsulting tendinum singultus

Modifications of Inflammation

Divided by pathologists into acute & chronic inflammation

Chronic inflam. It is generally attended with an increased secretions from the parts

It is frequently the result of acute inflammation

We are the creatures of habit. Habit is the cause of many chronic inflammations. Hence to cure them we must substitute be regular in our habits

Treatment of ^{acute} Inflammation

A suitable regimen is 1 of the most effective modes of treating inflammation consists in a proper confinement. Suitable diet

Aliphlogistic articles of diet consist of decoctions & infusions of vegetable substances as Rice water, Barley water, bread water, gum water, Apple water &c. & of sago, tapioca, arrow root

Active Treat

Blood letting next mode of treating ^{acute} local inflammation

Pulse is our best guide

Frequency of pulse ^{in health} is the number of beats in a given time

during 1st year 110 beats in a minute 2nd year 90 to 100

3rd year from 100 to 108 70 to 75 is average frequency of pulse in adult

Pulse is increased 5 beats in a minute on being raised from a horizontal posture

10 Modifications of pulse

1st Strong 2nd weak

3rd frequent 4th slow 5th hard & 6th a soft pulse

7th a full pulse & 8th a small pulse 9th an irregular or intermitting pulse & 10th an irritable pulse

Inflammation

c No 19th

1st Strong pulse if no other modifications of pulse does ~~it~~ not occur is no indication of disease

(It is found in individuals of a robust & plethoric habit ~~who~~ those who use plenty of exercise in open air

2 Weak pulse ^{more circumstances of weakness not to be regarded as indicating} occurs in individuals of delicate constitutions ^{disease} pulse may readily compressed which indicates the weakness weak pulse habitually of 120 indicates some organic disease

3 Frequent pulse when more than 85 When above 115 & patient is not owing to idiosyncrasy there is some organic affection

4 Slow pulse when less than 65 in erect & 60 in horizontal posture Unless owing to idiosyncrasy where it is habitual indicates disease almost invariably owing to some derangement in digestive organs Pulse is at once very seldom low & weak

5 Hard ^{or tense} pulse - corded pulse Indicative of some local disease Coat of arteries appears contracted It is mostly more frequent than natural but seldom above 115 mostly from 90 to 115

6 Soft pulse when of natural frequency is pulse of health

7 Small or thready pulse Not an uncommon modification of pulse It is to be taken as an indication of disease Artery appears to be constricted in calibre & feels thready It is most frequently in acute inflammation of serous membranes Generally of peritoneum

8 Irregular or intermittent pulse Very often arises from disease in or about the heart It is to be regarded as an indication of some local disease going on in some part

9 Irritable pulse occurs in the progress of many acute diseases It is probably always more frequent than in a state of health It is hard or utterly impossible to distinguish it from a hard bounding pulse without previous history of case It is frequent in hemorrhages also in blood letting where it has been carried to two great an extent

* 10th A full pulse used in contra distinction to a small pulse

Blood letting in treatment of Acute Local inflammation

It is the most powerful remedy in the treatment

Mode of performing the operation of blood letting

Should have all things ready - basin 2 or 3 cups a couple of bandages piece of lint Glass of water or a little vinegar & hartshorn 20 or 30 oz blood is a full bleeding 40 oz is a large blood letting for a stout young man

Inflammation

Effects of cathartics in treatment of acute local inflammation

besides their evacuant effects they produce revulsive applications
The kinds to be employed There are 3 kinds of purgative cathartics
Laxatives manna, magnesias, Elixir Pectinatum, Super-tartrate of
potash, aloes, sulphur black mass

Purgatives C. Galapae. C. scammonia Ol. Ricinae S. Soda. S. Magnesia

As a general rule these are the best in treat of inflammation
Drastic cathartics Gamboge, Croton oil, Nardica Elettarium white
& black helatone Ceyanthidis pulpa They are used only when
we wish to produce very powerful effects

Antimonial in treatment of acute local inflammation Their most
obvious effects is to produce diaphoresis

Tartrate of Antimony is best preparation of antimony
Pulvis Antimonialis or James' Powder when properly prepared
it is a most excellent preparation

It is seldom found good as obtained from shops

Relief in treat of Local inflammation

Blisters are undoubtedly the best They (blisters) have a stimulating
effect on the organs of organic life

Rheumatism

is divided into the Acute & Chronic
form Acute Rheumatism

fever gen. precedes the inflammation pulse is small, hard, & frequent
at commencement when fever 1st begins there is wandering
pain ~~over~~ after a while the pain becomes fixed It generally
attacks 1 of large joints Fibrous textures is the parts affected

Sometimes it involves the whole of the muscles of ~~organic~~ life

Characters of pulse alter pulse becomes bounding, full, & hard

Blood is of a highly inflammatory character urine is always of a
high colour Sometimes the inflam. takes place before the fever

It sometimes extends to the organs of organic life by metastasis
Terminations It has no determinate course Generally terminates
by a gelatinous effusion

Causes By far the greater number of cases arise from cold & moisture
There is a predisposition existing in some individuals which is gen-
erally hereditary

November 30th Acute Rheumatism

Most characteristic symptoms that it attacks the fibrous textures
It most frequently occurs in middle period of life & in
winter than spring

Treatment It is a sthenic or phlogistic disease
Blood letting is in the greater majority of cases proper
In common we ought to bleed freely & fairly

Purgatives It should operate freely - strong infusion of senna
with cardamom seeds to which may be added a little *S. magnesia*

Diaphoretics Spirits of Menerri may be giv freely as a
diaphoretic

Colchicum Autumnale is a valuable remedy It may be giv
in form of vinous infusion of seeds

Formula	{	Sulp. magnesia	23
		Calcinate	1-
		Water	13

vinous infusion 20 to 30 gr

It produces diaphoresis & keeps bowels open

Gallate of veratrin is the active principle of *Colchicum*

$\frac{1}{4}$ to $\frac{1}{2}$ gr is equal to 20 or 30 drops of vinous infusion

Many of organic affections of heart are caused by translation of Rheumatism

Opium nothing is to be gained by giving opium in any form in early part
of treatment until you have depleted fairly When giv the
dose should be large = 4 or 5 gr - combined with calomel

Chronic Rheumatism Generally the sequelae of acute stage

All the symp. exist in a less severe degree than in the acute form

It attacks itself to some particular part from which it is with difficulty
removed when you can remove it you may very often effect a
cure Paralysis from rheumatism very seldom remains long

Treatment Remedies are generally essentially diff from those used in acute form
Local blood letting sometimes required

Remedies internal & external 1st Guaiacum is a pungent stimulating
aromatic It may be called a laxative Aromatic tincture of Guaiac
is 1 of best remedies It should be giv with milk in as large doses
as stomach can bear

Oleum Terbinthina still better remedy Best remedy I have ever employed
should be giv pretty freely In large doses it acts on bowels instead of kidneys
It may be giv in rheumatism even where it partakes of acute form

December 1st

Rheumatism

1835

Formula { Gum Gutturina 13 }
 { Honey 33 } Table spoonfull
 { Water 53 } 3 or 4 times daily

When it is brought on by exposure after ptyalism Decoction of Sarsaparilla
 in large doses have failed

Calomel so as to produce ptyalism sometimes removes it after the remedies
 It should be giv. at night

Formula { Calomel 40 gr }
 { Opium 20 gr } 20 parts 1 every
 { T. Emetic 5 gr } night

External Applications When it attacks deep seated parts
 should use Cupping connected ^{followed by counter} with counter irritation alternating
 them successively Plaster of Pix Burgundica in lumbago

Simple Inflammation of

1st Simple Inflammation of Different textures of brain

1/2 of blood sent out from heart goes to brain

The pulse is not to be taken as a guide to inflammation of brain

In present state of our knowledge we cannot tell what texture
 of brain is affected Inflammation is apt to travel from
 1 texture to a diff. texture

Symptoms are same whether it exists in membranes or parenchyma

They are of most frequent occurrence in childhood

Cerebritis Inflammation occurring within cranium

Modifications of inflammation

1st Thenceitis Idiopathic It is of very rare occurrence

Most common causes Accidents inflicted on head Heat of sun
 irritating applications metastasis Alcoholic liquors

Characterized by Fever & Delirium & generally terminating in 4 or 5 days

Symptoms Loss of appetite Thirst, disturbed sleep, severe symptoms

fever patient feels as if a band of iron was tied around

the head ringing in ears in 1st period eye is dull & heavy

in 2nd stage eye becomes brilliant & sparkling 4th symptomatic

fever increases pulse increases Spontaneous vomiting takes place

bowels are uniformly constipated so that it is necessary to have recourse

to drastic cathartics Intellectual functions in progress become

furious There is now furious delirium By 4th or 5th day fever

begins to subside Evidently connected with great exhaustion

Cerebritis

Dec 8th 1835

Patient has a profound sleep pulse sometimes becomes natural
They denote that decomposition is at hand paralysis of 1 side
generally preceds death

Meningitis or Hydrocephalus It occurs at all periods of life

It 1st puts on a form that is not at all alarming It most frequently
occurs in childhood especially while child is cutting teeth
Appearances of stools They resemble parsley generally relieved by ^{1st visit by} giving
a purgative & by scarifying the gums sometimes however it continues
Epidemic frequently occurs child sleeps almost continually
1st serious cause of alarm is spontaneous vomiting Perhaps it will
occur once an hour even while child sleeping without its waking
It consists of bile There is no longer any doubt that inflammation of brain
exists Dilatation of pupil is a frequent symptom Sometimes the
symptoms are very insidious & are hardly observed by a common observer
as being any thing serious & Convulsions Coma & death generally follow
you are not always to find evidences of disorganization

Another Modification of Cerebritis very frequently found in young children
Sometimes it is congenital & sometimes the result of accident

It is of a more ^{chronic} character than former variety affection occurs gradually
3rd Form of Cerebritis arising from purulent discharges from ear
One of the most common sources of inflammation of brain

4th Modification Softening of brain (Ramollissement du cerveau)

5th Modification of inflammation of brain connected with tubercles They are
generally ^{found} in persons of a scrofulous constitution Frequently connected
with tubercles of lungs Diff in size from size of a millet seed to that
of a pullets egg This modification of Cerebritis is often of long duration

It usually terminates with symptoms of acute inflammation

Prognosis In all the modifications of cerebritis are unfavorable

If we can detect the disease in the early stages & adopt & energetic treatment
we may cure a great number

Diagnosis headache when complicated with other symptoms as vertigo

Intolerance of light & sound vomiting Spontaneous bilious vomiting
pulse is no diagnostic symp of itself It ranges from 50 to 120 in
course of a few days character of stools not to be relied upon generally
resemble parsley Coma convulsions & paralysis when other symp exist
are the most diagnostic signs

Cerebritis Dec - 10th 1835

Treat. of inflam of brain When you have satisfied yourself that cerebritis exists, you must use the most highly antiphlogistic treatment you should bleed to the greatest extent. Some recommend opening the jugulars when it has been carried as far as allowable then should bleed locally. Cathartics. Drastic cathartics should be had recourse to when stomach is not too irritable. In this case we should give large doses of calomel say 20 gr. When the irritation is relieved should give senna tea. Scammony in less irritable states of stomach may be given in combination with calomel. When these means fail should use croton oil in dose of from 3 to 5 drops. Cold applications. Most convenient mode is to take an ox bladder partly filled with iced water. More powerful mode is to apply cold water by means of a napkin. Most powerful effects can be produced by pouring from a spout. Among local applications are blisters very valuable in latter stages. Mercury carried to extent of inducing Syphilism. When given in large doses it keeps up purging. It is here only that it is useful. Tartrate of Antimony where it can be borne given in large doses it is a very valuable remedy.

Rachialgitis Inflammation within spinal cavity. I mean to include under this term all the modifications of inflammation of spinal marrow may be affected with all those morbid affections which have been observed in the brain. I shall consider all the Inflammatory affections under 3 heads. 1st Inflammatory affections where the seat of the disease is shown by the symptoms & ^{not morbid examinations} Symptoms of inflammation of medulla or meninges occurs in every age & sex both sexes more frequently in females & generally after the age of puberty. There is a morbid sensation in the part under pressure. The pains will shoot around the body in direction of the nerves. If it occur ⁱⁿ about the dorsal vertebra there will be morbid sensations in the epigastrium. In some instances the whole of the vertebral column will be affected. It will cause violent spasms in all the voluntary muscles. Sometimes there is a complete loss of both sensation & voluntary motion. In cases of this kind there is more or less symptomatic fever. In violent cases the bowels are obstinately constipated as a general rule. Causes. Sometimes the result of cold. Injuring &c

Rachialgitis

84

December 17th 1835

2nd Clasp. That is where the symptoms exist but where we have no demonstrative proof on account of patients getting well. If there is morbid sensibility in the part under pressure together with disordered functions of the cord you may be sure that inflammation exists in the part. To make pressure you should begin at upper vertebrae. The neurosis as chorea tetanus &c have their seat in medulla spinalis.

Dorsal vertebra especially 3rd & 4th are often found tender under pressure ^(in hysteria)
Pain generally shoots around to left mamma.
Treatment of inflam. of medulla spinalis. The remedies are few, simple & obvious. In a great proportion of cases the disease is local in which case the treatment should be local.

General rem. are required when constitutional symptoms exist.
Local Blood letting is generally required. Indeed it ~~is~~ ^{is} the indispensable counter irritation. Master of Burgundy pitch sprinkled over with Iodine is the best. It is in the main quite adequate. If these fail moxa & the caustic issues should be tried.

Sometimes it is more effectual to apply cupping to sides of spine while counter irritating applications are made to the other. Among other remedies it is sometimes necessary to confine patient to horizontal posture. Patient should be restricted to the strictest ~~to the~~ ^{to the} hygienic diet & regimen. In some cases 3 doses of the carbonate of iron is very useful.

I now proceed to speak of some of the Neurosis
Hysteria Paroxysm of Hysteria. Sometimes it comes on in a sudden & unexpected manner. Hysterical symptoms generally precede. Patient will weep & cry for a time then perhaps she will burst out into a fit of laughter. They frequently alternate. Another symptom is globus hystericus after it appears the patient will have a fit. It occurs in females most frequently between the ages of 20 & 45. It generally terminates by a copious discharge of limpid urine. It most frequently occurs in delicate females. It is generally connected with some derangement of catamenial discharge. Sometimes it puts on the character of a fit of apoplexy. Most frequently however it assumes the character of a convulsion. It sometimes becomes of a chronic character. There is gen. pain of left side. It is gen. connected with tenderness of dorsal vertebra with pain shooting around into breast. There is frequently palpitatio of heart.

Neurosis

Hysteria

Decem 15th 1835

Treatment Where paroxysm is violent if there is general excitement

should resort to a moderate bleeding & then give an emetic
 We must tend to the state of the spine, to the digestive organs & to the functions of
 Physicians should not show any alarm in treat of a paroxysm as the patient
 is sometimes alive to his behaviour in some cases

Chorea Sancti Viti nearly resembles hysteria Distinctive character
 is a twitching of some of the voluntary muscles

Epilepsy

Treatment strictest regimen & diet & vegetable diet Nitrate of
 silver Spirits of turpentine very valuable remedy

Cataplexy

Phlegmasia of the Respiratory Organs

Decem 22nd 1835

Catarrh 2 varieties catarrhus simplex & Catarrhus

Epidemicus

1st Catarrhus Simplex It is the mildest form of inflammation in the parts
 It may often be cured by giving even the most diffusible stimuli
 hence the adage stuff a cold & starve a fever

Most common exciting cause is changes of temperature

When the functions of an organ are increased the more liable it is to disease

Catarrhus Epidemicus or Influenza It most frequently occurs in cold &
 variable seasons It occasionally puts on a very favourable character

It seems to be brought about by certain morbid causes that exist in the atmosphere

Symptoms They are the same as in simple catarrh only more violent

Most inclined to attack middle aged persons

Treatment In the milder cases of catarrh very little is to be done
 Person should avoid exposure take less food than natural & use some
 mild purgative Bloodletting is a valuable remedy in some epidemic
 catarrhs Gen all that is required is to avoid the exciting cause
 use some mild purgative

Palati Tonsillitis Cynanche Tonsillitis

Symptoms difficulty in swallowing owing to the stiffened & enlarged

Tonsils voice becomes altered It is by no means an alarming

disease Sometimes the parts suppurate It is sometimes propagated
 to the cellular membrane so that the parts become very much swelled - lives

Treatment In mild cases all that is necessary is to avoid exposure use an
 antiphlogistic diet & a purgative medicine In violent cases decided treat is required

Decem. 29th 1835

Laryngitis

Cynanche Laryngea

I shall consider Laryngitis under 2 distinct heads 1st as it appears in children 2nd as it appears in adults although they occur in same structures yet they diff in some particulars 1st owing to the different degrees of development of Larynx

Causes It most frequently occurs in winter & spring & in low & damp situations

Symptoms It most frequently occurs in night Cough is of a very peculiar character It has been compared to the barking of a young dog voice indicates the same degree of hoarseness as the cough

In early periods of disease it is almost entirely local

It comes on gradually Febrile symptoms gradually increase Inflammation is seldom confined entirely to the Larynx Ordinary period occupied by disease is from 2 to 5 days

It is obviously divisible into 3 stages

1st The forming stage that is where the disease is purely local no symptomatic fever existing 2nd Stage commences with the symptomatic fever

3rd Stage where there is a formation of false membrane

1st Stage characterized by absence of fever, a peculiar cough difficulty of breathing & a slight hoarseness of the voice { is present

2nd stage where all former symptoms in a more violent degree & symptomatic fever

3rd Stage where we find respiration crouping & stridulous with a sensation as if the patient was breathing through a narrow tube & countenance becoming purple it indicates the formation of false membrane

Diagnosis It is a strongly marked disease generally easily to be distinguished It is apt to become complicated with measles

Treat of Croup It is very apparent

If we treat it in the 1st stage we may almost certainly cure it

1st remedy Emetics in early stages watery solution of tartaric acid 2 gr may be giv. to a child at a dose Should use the strictest antiphlogistic diet May give calomel in dose of 8 or 10 gr to evacuate bowels if Emetic does not produce that effect in 2 hours Having produced cathartic effects should keep down phlogistic state of system 30 or 40 d of a solution of tartaric acid in proportions 4 gr to oz water Dr Rush seldom found it necessary to bleed in this stage

2nd stage When called to a case in 2nd stage & emetic is not the 1st remedy Blood letting should precede all other remedies Sometimes do in case of very young children it becomes necessary to bleed from external jugular As a gen. rule 3 or 4 oz of blood should be taken from a child 2 or 3 yrs old Blister, warm bath, Calomel - It should not be relied upon as a main remedy

Laryngitis

December 30th 1835

3rd Stage occasionally leeches, warm bath Polygala senega. Last resort is tracheotomy which holds out only a faint hope

Laryngitis as it occurs in Adults. It is of rare occurrence

It does not gen. begin like croup in children. It gen. begins like palato tonsillitis gradually passing downwards into Larynx. Cough is of a sharp whistling sound. It is generally attended with severe pain about thyroid cartilage. Generally speaking it terminates about 2nd day. It seldom extends beyond the larynx.

Laryngitis in adult is more violent & dangerous than croup in child.

Treatment great principle is same as in the treatment ^{of croup} as it occurs in children.

Blood letting generally & locally. Nauseating doses of tartar emetic blisters &c. Bronchoderm, is more likely to succeed in Laryngitis of adults than children because in these the inflammation generally extends through the ramifications of the bronchia.

Pneumonia of the textures of the organs contained within the Thorax

Acute Bronchitis

Inflammation of mucous membrane lining ^{trachea} ~~bronchia~~

It is one of the most common diseases in all parts of N. States occurs most frequently in winter & Spring.

It usually begins like a common catarrh in fact it is often brought on by frequent attacks of catarrhs.

Cough comes on in paroxysms. There is generally some soreness felt on coughing. Cough is wheezing. Whenever patient is exposed sudden change of temperature a fit of coughing comes on. Sputa is gen. thin & acrid. One of most frequent symptoms is pain in head especially when a fit of coughing comes on.

There is uniformly disordered functions of digestive organs.

Urine is gen. turbid small in quantity & high coloured. Pulse is small & frequent. Though high inflammatory symptoms are present as flushed face &c.

There is an uneasiness gen. felt about precordial region but no distinct pain. Patient generally has a fit of coughing on going to bed. Can lay on either side.

About 4 or 5th day about 4th or 5th day expectoration changes & becomes more copious which if it continues to increase becomes unfavourable.

In favourable cases when expectoration becomes free the symptoms begin to abate. When inflammatory symptoms are most severe the pulse is small & frequent.

A moderate general bloodletting in early stage is beneficial.

88

Bronchitis Jan. 5th 1836

Acute Bronchitis characterized by great oppression & tightness in the breast headache greatly aggravated on coughing General prostration of ^{system}
 Treatment It requires great delicacy of management Disease is highly phlogistic General Bloodletting ought to be practiced though with great caution & only in early stage After expectoration takes place gen. bloodletting ought not to be employed Should bleed from a small orifice & only to the extent of 10 or 12 $\frac{3}{4}$

Antimoniala are among the most useful remedies It may be carried to the extent of inducing vomiting At same time should allow the free use of demulcent drinks After expectoration has occurred tartar E. should no longer be used Emetics are however required occasionally in children - Squills Cathartics cannot be used as freely in pulmonary as other diseases Milder purgatives best Blistering is a valuable means Opium Injurious in early stages It may be used after expectoration has taken place The mind becomes exalted as the corporeal frame becomes weakened Diet & regimen They must be strictly antiphlogistic

Chronic Bronchitis

Following is gen. its course

Patient has a cough especially in morning on waking Expectoration is viscid & yellowish It is of a globular form Patient eats a moderate breakfast at dinner appetite is capricious In afternoon patient becomes weary Patient is very sensitive to changes of air patient becomes emaciated pulse feeble & frequent The mind becomes exalted as the ^{corporeal} powers become weakened most prominent symptoms cough & copious expectoration Ectetic fever generally supervenes

Post mortem appearances The lungs do not collapse

Treatment General Bloodletting Very few cases that require it

It being generally attended with great debility, only cases which it is where there is hemorrhage from lungs

Local Bloodletting useful in treatment of chronic bronchitis

Counter irritation Blisters, setons, & issues &c

Emetics are occasionally required In Specac. 1 gr Speace 5 gr powder ^{Emetic}

Expectorants not beneficial Wine of seeds of colchicum useful rem

Balsam Copiva I have never seen it do good in treat of C. B.

Mercury so as to produce slight salivation is a very important remedy

Bronchitis

Jan 6th 1835

Mercury It should be given in small doses

Formula

Calomel 12 gr
Opium 12 gr
Tquill 8
Antimony 2 gr
divide into 12 pills

Take a pill at night on going to bed

Calomel is the medicine from which the pill receives its efficacy

Sulphate of Quinine sometimes useful 1 gr with $\frac{1}{4}$ to $\frac{1}{2}$ gr
Opium may be given 3 or 4 t. d.

Night sweats best relieved by Acidum Sulfuricum Aromaticum
Diet & Regimen Diet should consist of farinaceous articles & milk
Drink, water Should guard against changes of temperature

Pneumonitis Inflammation of the parenchyma of Lungs

Acute & Chronic

1st Acute Pneumonitis It comes on with a smart symptomatic fever accompanied with pain in some part of chest Respiration is short & hurried attended with a sense of anxiety about precordia Patient is but little inclined to indulge in speaking as it increases the pain Respiration is increased in diff so that the patient is unable to lie down Face is pale & eyes suffused

Expectoration at 1st is scanty It is bloody expectoration is uniformly of a red colour - most pathognomonic symptom In advanced stage expectoration is only striated with blood Pulse is excited It is frequent, hard, & tense

Tongue is covered with a white fur There is complete inappetency, thirst

Blood is uniformly cupped & buffy Delirium occurring is an unfavourable symptom Percussion & mediate & mediate auscultation are a valuable means of examination Hepatization results from a deposition of coagulating lymph

When both lungs are involved the case is rapid in its progress

It may terminate Hepatization odema Suppuration or gangrene

Causes - It may occur at any period of life but more freq. after puberty up to 45 Those persons who inhale irritating substances into lungs are liable to it

Most frequent in cold damp weather & winter & spring

measles freq. fol by Pneumonitis It has occurred as often as 20 times in some individuals

Treatment Blood letting always to be practiced in its treatment

Should bleed from large vessels & carry it away to syncope Should be

fol. by some purgative Sometimes necessary to bleed a 2nd time in course of 10 or 12

Antimony very valuable in treat 1 gr ev. 2 hr increasing the dose

Blisters &c

* Most characteristic symptoms Fixed, obtuse, circumscribed pain expectoration being intimately mixed with blood

Chronic Pneumonitis sequelae of catarrh & Pneumonitis &c

In *A. Pneumonitis* where patient appears convalescent but the pulse continuing frequent say from 90 to 100 it indicates a termination in chronic *Pneumonitis*.
 Mind is often as much exalted as in chronic *Bronchitis*.
 Cough is freq. attended with mucus expectorations mixed with blood.
 Patient becomes emaciated. Percussion & Auscultation are valuable means in forming a correct diagnosis. It is manifestly owing to hepatization of lungs when unconnected with tubercles it is not necessarily fatal. We may fairly hope to save the patient. When by percussion & auscultation we discover the locality we should cup over the part also use counter irritation over part as blisters, pustulations &c.
 Mercury so as to produce slight ptyalism very beneficial.
 Diet & regimen. Patient should live upon a vegetable & milk diet. Should avoid exposure & also use exercise in carriage.

Pleuritis January 7th 1830

Acute & Chronic — *Acute Pleuritis* freq. connected with *Pneumonitis* & *Bronchitis*. Symptoms are generally strongly marked.
 A severe pain gen. confined to 1 side of lungs. It mostly occurs in lower portions of lungs. Pain much aggravated on taking a full inspiration V & V.
 Expectoration has nothing remarkable about it. It occurs most freq. in persons with narrow chests & in those liable to catarrh (*Bronchitis*).
 One attack seems to be somewhat a preventative of a 2nd attack.
Treat. In regard to treat. very little reg. to be said. Blood letting should be to as great an extent as in the treat. of *A. Pneumonitis*. Emetics should be freely used. Replication of great importance.
Chronic Bronchitis Pleuritis the vital energies soon become broken down by frequent relapses. Effusion occurs. Countenance becomes bloated & livid. Patient cannot lie in horizontal posture. When instead of serous effusion matter becomes effused the symp. are all aggravated. { chest
Treat. It does not differ essentially in treatment from other chronic inflammations within

Phthisis Pulmonalis It is not to be considered as 1 of the Phlegmasiae

Tubercles 2 most general forms which they assume 1st Where they exist in distinct granular masses. They seldom exceed the size of a cherry stone. When they have arrived at this size they become ^{softened} in centre & continue to soften until they become completely softened & are then expectorated. Tubercles develop themselves most early in upper part of lungs. This is not at all uncommon. 10th find excoriation in lungs so large that you can introduce whole fist. Tubercles are the products of inflamma

Phthisis Pulmonalis that class of chronic cases attended with cough emaciation hectic fever &c & upon examination after death tubercles are found in the lungs

Prominent distinctive features of the disease

Cough generally speaking cough is the most troublesome symptom Sometimes it is almost wholly absent

Expectoration at 1st scanty At later period of disease it increases in quant & becomes of yellow colour owing to tubercles being effluorated

Hemoptysis is the expectoration of a large quant. of blood Sometimes it is small in quant. & is coughed up with the mucus

It is a strong indication of *Phthisis Pulmonalis*

Pain not so an invariable a symp. as is supposed by many When it does occur it is by tubercles irritating pleura

Hectic fever is the most uniform symp. of *Phthisis*

Another symptom is lesion of stomach occurring in course of disease The stomach becomes especially painful Diarrhea is almost an invariable symptom in latter stages of *Phthisis*

Diagnosis before stethoscope had come into use *its* diagnosis was inconclusive There is no dependence to be put in chemical or other tests Vomices are nothing more than large tubercular masses which some times burst & become expectorated Pieces of this kind we may be sure that tubercles exist in lungs

Galloping Consumption owing to a large number of tubercles being developed at once so that expectoration of tubercles takes place so profusely that the patient is destroyed in a short time The lungs being unable to perform ^{their office}

Stethoscope is almost invaluable assistant in forming our diagnosis (the question)

The hope of affecting a cure is in the present state of our knowledge is out of the question

Laryngeal Phthisis It is an inflammation of mucous membrane of larynx attended with a thickened condition of its structure There exists an irritation about the Larynx with cough - difficulty of speech sometimes a complete loss of voice Difficult expectoration owing to mucus about Larynx being affected

Causes of *Phthisis Pulmonalis* Sometimes acquired

It recurs more frequently in women than men It is freq. transmitted from parent to offspring Cold & damp climate mechanical irritations

Treatment of Phthisis Pulmonalis Jan 1836

Bleeding may sometimes be employed as in Hemoptysis As a general rule it is ^{improper} Sulfate of Copper so as to produce nausea has been recommended where Hemoptysis occurs also sugar of lead Nitrate of Potash &c The tendency of these remedies is to depress the system when tubercles are beginning to be expectorated there is a danger of system being too much depressed

Counter Irritations Letons, Issues, Moxa, actual cautery very useful remedies where there is ^{local} inflammation existing

Among the remedies spoken of as specifics are

Digitalis Purpurea It is worthless in treatment of Phthisis

Mercurial remedies their use instead of curing increases the development of ^{tubercles} Prussic Sea air Chlorine & Sodium gases they all fail to be of any use

Plegmasia of the organs contained within cavity

of Thorax & Abdomen Jan 18th 1836

Gastritis or inflammation of mucous membrane of stomach

Frequently occurs in connection with aphthous affections of mouth

Symptoms They are not striking A sense of pain & heat about epigastric region appetite is impaired generally lost Thirst tongue covered with a thick fur with a red apex & edges Great pain often felt after taking cold drinks almost every thing that is taken into stomach is vomited up pulse is frequent & small countenance is haggard & anxious mind querulous As an Idiopathic disease it is uncommon 1 in 100 patients

In progress of many acute diseases the stomach of law becomes inflamed occasionally met with in all ages in all both sexes & in every climate but most frequently in hot seasons & in hot climates

All the organs are in a state of increased state of excitement in hot weather

Chronic Gastritis frequently conformed with dyspepsia

Treatment Expectant method chiefly to be relied upon as all drugs have a tendency to produce irritation

In selection of diet should chose those remedies that are antiphlogistic & soothing as vegetable decoctions & infusions Barley water rice water acidulated with lemon juice miscelages of gum arabic &c Gums are the most strictly antiphlogistic & at same time the most soothing &c

Emetics should be given together with some laxative medicine as calomel

Most common symp. are pain & morbid symptoms sensibility in Epigastric Loss of appetite nausea & vomiting

Blood Letting General Blood letting sometimes required.

Local Bloodletting as there is not much symptomatic fever is gen. preferable
Counter Irritation very valuable as dry cupping Sympiesms
blisters &c

Emetic a mild emetic where there are symptoms of impurities in stomach
copious draughts of aniseed tea

Inflammation of Intestinal Canal Jan 20th 1830
Dysentery Character Causes Symptomatic Terminations
Prognosis Treatment It is a phlogistic disease

Blood letting should be employed freely It is not a disease which we should expect
to cure by a copious bleeding but it gives efficacy to other remedies
Cupping also a valuable remedy

Cathartics Castor oil best purgative Next best is sulphate of potash
In dose of ʒij every 4 or 5 hours until it purges

Mercury so as to produce slight Stomatitis is a valuable rem. in treat of Acute
dysentery especially of warm climates

Rest in horizontal posture should be rigidly enjoined

Abstinence all important demulcent drinks may be freely allowed

Opium when restored to it should be given in large doses 2 or 3 gr doses

Emetics in some cases very valuable Blisters are of no use

Cholera Morbus

It differs essentially from Cholera Asphyxia It occurs in every part of U.S.
but most frequently towards South & West Generally most ripe in August &
September when nights become cool

Symptoms 1st place it generally occurs at night & quite unexpectedly to patient
Individual goes to bed without indisposition & is awakened up with derangement of digestive
apparatus such as nausea with a disposition to go to stool There is a feculent evacuation
of feculent matter from bowels Vomiting takes place in a short time Discharges
continue to increase Stools after 1st are very watery

Pulse is seldom excited in any period of the disease

Treatment It is in some rare instances connected with local inflammation ^{which will modify treat.}

1st object ^{is to stop} the morbid secretions & soothe the system If any undigested food
remains we should remove the irritating cause effectually before we use any other treatment
by giving a table spoonful of corn salt or a little mustard

Opium It may be given in the form of black drop in large doses 30 or even 40 drops
on a little sugar Sometimes it is best given in solid form to prevent it from being
thrown up & patient should not be allowed any drink for a time

Cholera Morbus Jan. 22nd 1836

Arresting Applications Sinapecs best Heat should be applied to extremities & body wrapped up in warm blankets Should not leave the patient as the time of action is short When opium is used freely it seems to act almost like a specific It is the chief & great resource for any of the diffusable ^{stimuli} Having stopped the discharge little remains to be done Allow the patient only a small quant. of drink at a time In 6 or 8 h after vomiting & purging you may give a dose of Calomel or Castor oil will generally succeed in producing copious ^(discharge) feculent

Cholera Asphyxia

History of the origin of Cholera It originated at Gashaw on the Ganges near Calcutta in India in the month of August 1817

It arrived at Moscow in 1830 Reached Quebec in the month of June 1832 60,000,000 of people or upwards are supposed to have died off this disease before it reached the U.S.

Premontory Symptoms as they occurred in N. S.

Stomach & bowels are indisposed The vulgar would say "Do there is a rumbling of my guts" Diarrhea occurs very early in complaint generally vomiting also occurs early in complaint Disease tolerably manageable at this period of disease which may be considered as 1st stage

Sometimes the patient is seized at once without any premontory symptoms

2nd Stage may be supposed to begin when decided vomiting & purging occurs

After 2 or 3 discharges they resemble rice water Features shrink skin becomes of a leaden hue Extremities become cold copious perspiration breaks out over body pulse is soon lost at wrist One of the most characteristic symptoms is loss of elasticity of skin If you pinch it up it remains in this condition Ends of fingers of purple hue lips blue Tongue gen. clean It looks & feels like the tongue in a dead person Breath is cold Superadded are cramps of extremities & abdominal muscles Loss of voice

Mind is seldom much disturbed Considerable time before death the heart acts very feebly Lungs have lost their function Bile, Urine & even tears cease to be secreted From commencement the thirst is intolerable Epigastrium hot when drink is given it is almost immediately rejected It is properly called the stage of collapse

Among the remarkable symptoms is a change of the voice

3rd Stage If patient survives 2nd stage or the stage of collapse the patient may recover The pulse returns at wrist discharges become mixed with bile Urine begins to be secreted 2nd attack seldom occurs in same epidemic

Cholera Asphyxia Jan. 25th 1836

Causes We know very little about the cause of disease & especially cholera
 3 opinions prevail as to cause of cholera 1st Attributed to some peculiar change
 in the electrical condition of the atmosphere 2nd To Contagion
 & 3rd To some unknown substance floating about in atmosphere

In some way or other it does possess the power of multiplying itself

Treatment of Cholera & considered under 4 stages 1st Premonitory 2nd Stage when vomiting & purging takes place 3rd Stage of collapse & 4th Stage of reaction

Premonitory Stage If taken at this period there is nothing alarming
 The remedies are of the simplest kind & supposing there is nothing but 2 or 3 fecal
 evacuations Should remain quiet in horizontal position Live on animal diet
 take a gr of calomel in combination with a gr of Elixar or 4 or 5 gr blue pill
 so as to alter the evacuations from bowels Such simple remedies will at once restore the
 patient to health A few drops of Camphor will almost constantly relieve the rumbling
 noise of stomach & crivels

2nd Stage 1st thing perhaps that is taken notice of is nausea & vomiting & purging
 The hazardous period has begun Blood letting decidedly pernicious in treat.
 of cholera as it occurred in Phila. & use of opium & the diffusible stimuli
 have been gradually laid aside They are now restored to in a more measured form
 Calomel

3rd Stage Treat the Treat should be
 When rice water discharges have commenced 1st remedy one of the milder emetics
 as mustard or cam. salt. Sinapisms to epigastrium Warm applications
 to preserve the animal heat Irritability of stomach having been somewhat allayed
 may give calomel

Formula Calomel 1/3
 Opium 12 gr } divide into 12 pills &
 Camphor 12 gr } give 1 every hour

If the case still grows worse you should change the treatment

Chlorate of Potash 8 gr
 Supercarbonate of Soda 1/3
 Cam. salt 1/3 scruple } dose 1 spoon full
 Cepid water 4/3 } six hour

This mixture has the effect of checking vomiting in some instances

Another Formula also recommended as very beneficial

Super C. Potash
 gum arabic
 loaf sugar 1/3
 elixer water 8/3
 3 or 4 of oil left

Carbide of Antimony as a remedy not more successful than other remedies that have been tried

Cholera Asphyxia Jan 29th 1836Treatment of 3rd Stage or the stage of collapseArtificial heat. Some persons say the temperature of the body is below ^{that of} surrounding objects

Wrapping patient up in warm blankets also by applying hot bricks

or by means of hot steam

Great emphasis laid on stimulating frictions

more especially in N York

Emmetta consisting of saline resin or turpentine

Injections into the veins is the most ^{certain} remedy of bringing about reaction

It may be blood or salt & water

Dr G. Mc Clellan injects 72 3/4 water

with 3/4 salt into the veins of an old woman who was apparently in asthenic morbus & thereby brought about reaction

4th Stage or stage of reaction

The fever is of a typhoid character

A great number die after reaction has taken place

Fever February 1st 1836

Idiopathic fever Def: All those acute diseases in which there is a general disorder of all the functions of the body without any obvious local inflammation existing as a cause

Predisposing cause of Intermittent & Remittent fever

is Malaria = produced by the process of decomposition of organized bodies

What are the circumstances under which Intermittent & Remittent occur

They occur most frequently in low & marshy situations although some parts of the country the people are exempt from them though they are surrounded by low & marshy situations

These facts go to show that we have not arrived at the true cause of these diseases. It appears that something else is required besides the decomposition of animal & vegetable substances. The inferior

animals are not affected by these fevers

Physical Properties of Malaria

It is a ponderable body & exists in

an aeriform state. Next physical prop^y is that it is manifestly absorbed by

salt water

February 3rd 1836

only known predisposing cause of Intermittent &

Remittent fevers is malaria. They seldom occur on Eastern continent

Living body becomes accustomed to Malaria in some degree

youth or a robust constitution is no protection from miasmatic diseases

The time at which it produces its effects on system depends upon susceptibility of system & the concentration of the malaria

Next subject of inquiry Intermittent fever

97

Intermittent Fever February 3rd 1836

In 1st place it is emphatically a disease of whole system

Rigors, increased heat & increase in frequency of pulse is characteristic of disease

Various Chronic affections are the consequences of this disease

In 1st place one of the most remarkable characters of the disease is its periodicity Interruption is complete in many instances

It is divided by Pathologists into Simple & Malignant

Malignant form puts on very diff. aspects

Simple Intermittents are divided according to diff. periods at which they occur into Quotidian, Tertian & Quartan &c

Some writers assert that it occasionally arises from other causes than malaria

As worms They only act as occasional exciting cause when patient has been exposed to malaria Only known predisposing cause is malaria

They occur most freq. in spring & Autumn If they occur in spring they are called vernal if in Autumn they are called Autumnal agues

An ague occurring as early as July is generally mild Vernal agues are the mildest & if left to themselves have a tendency to get well with respect to paroxysm

usually divided into 3 stages viz cold, hot & sweating stages

Patient is attacked with heaviness & yawning Coldness generally begins about the back feels as if water was running down back Chilliness sometimes begins in extremities The nails appear purple

Chills appear to spread in waves over the body There is an appearance of goose flesh pain in various parts of body & great distress of mind

Shivering takes place sometimes very violent Cold stage generally lasts about 24 pulse is very much diminished in volume & force

Respiration is difficult there is a sensation of thirst with a burning heat internally There is pain extending around from spine to abdomen

Urine is small in quant & high coloured When cold stage is terminating the symp become mitigated The pulse now begins to act with greater force

Respiration becomes full vital energies begin to act with preternatural vigor skin becomes red the sensation of heat is restored The temperature

sometimes rises very high great thirst, throbbing of carotids In children epistaxis is very liable to occur during hot stage patient is very restless

lying about from side of bed to other

3rd Stage or Sweating Stage as soon as sweat breaks out the symp begin gradually to disappear pulse becomes natural

Intermittent Fever February 3rd 1835

The urine becomes large in quantity & deposits a catenitious sediment on standing. The patient is quite well during intermissions. Appetite is gen. somewhat impaired. Usual duration of paroxysm is from 5 to 8h. Intervals are generally free from fever though the patient experiences some degree of derangement in health. When the disease returns at an earlier period of the day than at former attacks it is said to be anticipating & is unfavorable. Postponing when it occurs at a later hour in day which is a favorable circumstance.

February 4th Quotidians are more unmanageable than Quartans. There are certain periodical diseases which have been called *males agues*. Certain other diseases occur periodically as Gout, Epilepsy, &c. Malignant Intermittents. They are exceedingly irregular. Sometimes reaction does not take place or the cold stage may last a great length of time. Comatose variety. The patient becomes drowsy.

Delirious form. Closely allied to the other forms. Next modification is the sweating malignant Intermittent. Sweat is poured out from every pore. The patient becomes cold. There is a remarkable tendency to fainting.

Peripneumonic variety. Paroxysm is usually very protracted. Gastric malignant. There is violent pain in stomach. There is a constant tendency to fainting. Intermittents occurring in aged persons dangerous. Gastric & Enteric affections are very common where the disease is protracted. Among the sequela none is more common than hepatic affections. Crimefaction of the liver & spleen are very common even in mild cases.

Relapses are very frequent in intermittents. No advantage is derived from allowing intermittents to run on a certain number of paroxysms. Epilepsy & cutaneous affections have been cured by them.

Post mortem appearances. Congestion of internal organs very common appearance.

Treatment of Intermittents. Something remarkable in regard to treatment. Some people suppose that the disease produces longevity. By others it is looked upon with absolute horror. It is cured by the most opposite treatment. With so many guides pointing out different modes of treatment it is hard to choose the right course to be pursued. Customs disarm a disease of its terror. In many cases they are mild & will get well of themselves.

Therapeutic Remedies are divided into general & specific.

1st direct your attention to the general state of the health.

Intermittent Fever February 5th 1836

Treatment Treatment during the paroxysm & also during the intermission
 During the Fit of simple intermittent very little requires to be done but it is otherwise in the malignant form. All the diffusible stimuli must be restored to

Treatment during intermissions Sometimes it is necessary to procure general bloodletting also recommended in cold stage Recently been revived by Dr Mackintosh I would use it in certain cases Most proper times is during intermission
 Opium has also been employed in cold stage
 It is undoubtedly a good remedy given in dose of 1 gr an hour before the return of paroxysm
 Emetics have also been advised before & also during the cold stage

It is a harsh remedy in cold stage Best used in cold climates
 They may be employed with advantage during intermissions Speac in combination with calomel is best (20 gr with 2 of calomel)
 Catharsis should be kept up during intermission
 With respect to the specific treatment

Peruvian Bark or Cinchona The most efficacious remedy we possess
 Dose 13 repeated as often as the stomach will bear

Sulphate of Quinine is generally to be preferred to any other preparation
 Best given in solution in water } 2 Quinine 16 gr } teaspoonful or 2 or 3
 When objected to by mouth it may } Sulph. Acid 16 gtt } each of which contains about
 be given in form of Emetic } Aq. font. 15 } 2 gr of Quinine

A new mode is to remove the skin & apply an ointment composed of 6 gr of Q. with 93 of lard
 Another preparation of Quinine is the Hydro ferri cyanate of Quinine Not aware that it can be procured in this country 2, 3 or even 8 gr may be given in course of 24 h
 Next specific is Fowler's solution of arseniate of Potash Dose for adult 10 d
 Mercury Sulphate Zinc Sulphur Cupri. Vascular Emetic Inferior tartrate of Potash
 Sudorifics as Decoctions of sarsaparilla &c

Remitting Fever February 6th 1836

In Northern latitudes The miasmatic diseases are gen. mild
 They put on their most malignant form in southern states It arises from same cause as intermittent fever Intermittents & Remittents are in some way or other produced by decomposition of animal & vegetable matter

They are divided into Simple & Malignant Remittents

1st Simple Remittent fever very frequently fatal

Simple Remittent Fever

February 9th 1836

It is the most common disease in this Country
 Treatment It is a highly phlogistic disease There exists a great
 tendency to the production of local inflammations in diff. organs
 Blood-letting is the chief anchor of our hopes It should always
 be employed It should be freely used

Should next evacuate the bowels When you have accomplished the evacuations
 of the bowels you have made a considerable stride towards effecting a cure
 Kind of purgative to be employed Calomel & jalap aa 10 gr - Rush
 I prefer calomel alone It will be retained better than any other medicine
 Should allow the patient very little drink at times To relieve thirst may hold
 a little cold water in mouth May give 4 or 5 gr every hour or two until 30 or
 40 gr have been taken Purgine not produced may give Sulphur Magnesia
 Having once succeeded in reducing the pulse & evacuating the bowels you should
 give small doses of calomel so as to produce slight ptyalism 4 or 5 gr w 3 or 5 h
 Acid Cathartics prejudicial Should use the strictest antiphlogistic diet & regimen
 Diet should be restricted to & also allow a little Seltzer water
 Bark should not be given until complete remission has taken place
 A few medicines as feasible should be used in the treatment

Malignant Bilious, or Yellow Fever Feb. 9th

It arises from modifications of same predisposing cause as Intermittent & Remittent
 Predisposing cause the same as in Intermittent & Remittent fever Malaria
 When it occurs early it is a sign that it will prove very fatal Sometimes it
 occurs as early as June seldom earlier than August When occurs Epidemically
 it is called yellow fever but when it attacks only a few individuals it is
 considered only as a severe attack of Bilious Remittent
 Prophylaxis I should make no essential change in habits

Treatment I consider it a phlogistic disease & use antiphlogistic
 means Should be governed by general principles as in treatment of Remittent
 Bloodletting is a most admirable remedy
 Purgatives of active kind are to be preferred Should obtain free discharges
 Calomel should constitute the principle ingredient & should be used freely
 fol. it up by giving Sulphur Magnesia
 Ptyalism after free depletion sometimes proves critical in the disease
 Emetics seldom used since the time of Dr. Rush
 Sometimes beneficial to use gentle emetic - Ipecac
 Cold affusions during the stage of reaction generally used in this Country by Sp
 Stimulation They will bear larger doses of Stimulents in this disease than any other

Doses of calomel 1836

101.

Continued Fever February 13th 1836

Distinguished from Intermitents & Remittents by their not originating from malaria & by their not coming on in paroxysms but being continuous. They continue for a longer or shorter time when they terminate in a crisis exacerbatation occurs about 7 P.M. in evening.

At commencement of a continued fever all the organs of the body are affected. They generally occur during the cold months.

Three modifications of fever Synocha Typhus & Typhoid.
Synocha an inflammatory fever Typhus is opposed to Synocha in all its forms.
1st Synocha in vascular tongue called inflammatory fever.

Sometimes the fever continues but Typhus is the very mildest of the idiopathic fevers.

Causes There does not appear to be any predisposing cause. It is found almost uniformly in robust persons. Any sudden shock given to the system may produce the disease. Atmospheric changes, hoarse, fear.

Found at all seasons and all climates. Most frequently in northern climates.

It will disappear almost as abruptly in some cases as it began.

Treatment Nothing peculiar about it. It is phlogistic & opposed by antiphlogistics. It is brought about by transient causes. Therefore nothing required but the mildest treatment. Treatment depends upon general principles.

Next of which I shall speak is Typhoid.

Predisposing Cause It is the most common of the infectious diseases.

It arises from a combination of accidental circumstances & also

It arises from persons labouring under the disease.

When individuals are confined in crowded & ill ventilated apartments persons going there are apt to contract the Typhoid fever.

Is it generated by the decomposition of animal substances. Effluvia arising from the human body is supposed to be more deleterious than that from other animals unless they are not in a very concentrated form. They are not injurious to health.

Where there is an infectious disease should use free ventilation avoid inhaling the breath of the patient & should keep out of the current of air as it comes from patient & give the attendants the same direction. Clothes about the patient should be frequently changed & immediately put in tub of cold water.

Fumigations vinegar is valuable Nitric acid Chlorine

Nitric acid for sick chamber is best should place it in diff parts of room.

Typhus Fever February 16th 1836

Characters of Typhus more especially a disease of cold climates
Not so common in this country as in the northern part of Europe though
it does occasionally occur in some.

Symptoms Classify it as you please but still the disease is typhus
There are 3 distinct stages 1 Stage of depression 2 stage of excitement
& 3rd stage of collapse

1st stage or stage of depression It for the most part approaches very gradually
1st general languor disinclination to move about The mind is generally stupified
as well as body countenance gen. pale haggard & dull slight chills in diff parts
of body digestive organs more or less impaired tongue shiny & relaxed
it receives the impression of teeth It gen continues from 2 to 4 days
2nd stage or stage of excitement In many instances the temperature rises
much above natural Appetite is entirely lost Inextinguishable thirst
an exacerbation takes place gen about 7 or 8 o'clock in evening & continues until morning
There is exacerbations & remissions ev 24h They continued with more or less
severity for 7 or 8 days when the stage of relapse or the
3rd stage occurs patient becomes much emaciated mortification apt to occur
in the parts upon which patients lie pulse becomes weak & very frequent
articulation becomes difficult & imperfect patient lies upon back & has a tendency
to slip down in bed Character of secretions about gums & lips They become of a dirty
yellow colour Teeth becomes sordid Sensorial functions The patient thinks
he sees something before his eyes superintention of twitching of the muscles is another
indication of great debility There is sometimes considerable difficulty of
swallowing fluids discharges take place unconsciously
fæstichia occurs They resemble a mahogany colour & are cov with a smooth thin
February 17th 1836 There is no disease in which there is so close an approximation
between life & death as in this & yet the patient recovers

In a majority of cases local inflammations does not exist
though in many cases symptoms of inflammation does exist in some of the organs
Brain is often the seat of inflammation Not so apt to occur in respiratory passages
It mostly attacks the parenchyma of lungs Local inflammations supervening on
typhus are dangerous Local inflammation of mucus membrane of stomach &c
of frequent occurrence Liver is occasionally the seat of inflammation though
rarely Sometimes congestion occurs in liver They very often terminate in inflammation
Typhus does undoubtedly sometimes exist & yet there is no local inflammation
in any course of disease

J. Brown

Post mortem appearances Most common appearance is ulceration of Glands of Peyer

Synochus Fever

I shall make only a few remarks upon it

It is ^{the} common fever of cold climates not however limited to cold climates
It is intermediate between Synocha & Typhus In common language it is
called typhus & also by many in the professions
I know of no line of demarcation only the mode of their origin by which I can be dis-
tinguished from the other

Genuine Typhus invariably arises from Infection { too exposed
Synochus arises from more accidental circumstances to which all are liable to
as Cold Moisture Debachery &c &c

Treatment of Synochus & Typhus

We can mitigate the symptoms of Typhus & curtail the disease

Its tendency is to break down the constitutional system

In 1st or oppressed stage our object is to bring about reaction

Should place the patient, in a spacious & well ventilated apartment, in bed

How shall we bring about reaction 2 modes 1st by diffusible stimuli

They are generally injurious Where it is excessive we are obliged to resort to them

An emetic is generally the best remedy - Tartar Emetic preferable either alone
or in combination with calomel It is desirable to produce catharsis at some
when prostration is not great

2nd stage or stage of Excitement There is a great tendency to congestion { inflammation

Blood letting is an appropriate remedy for treat of both Synochus & Typhus

when cautiously used & at proper period In advanced period of 2nd stage

it is necessary to bleed if inflammation occurs in an important organ

must proceed with great caution If brain becomes inflamed should also apply cold

Purgatives are almost invariably employed during stage of excitement

20 or 30 gr calomel fol. by comp. tinct senna

Cold Effusion It is one of the effective remedies Manner to be practiced

Patient taken from bed & cold water poured on patient from head to foot or

by wrapping patient up in wet blanket Avoid using it where there is local inflammation
of any extent Pulse falling irreg. heat reduced, patient falls to sleep &c

3rd stage, stage of collapse Indicated by falling of temperature &c

Sometimes impossible without trying the effect of stimuli to tell when it is proper that
they should be administered

Notes taken from Dr. J. Revere's

Nov 14th

Lectures on Practice of Medicine 1836

Inflammation

Nails, hair, epidermis, enamel
are the only parts which are not susceptible of inflam.
Distinctive characters of inflam - pains redness
heat & tumefaction

Redness

is always present in inflammation

Redness may or may or may not be an evidence
of inflam. after death

Heat

Patients in the article of death almost always be on back
It constitutes a remarkable phenomenon of inflam.
Is the inflamed part hotter than blood at heart

As far as direct experiment goes it is not

Tumefaction - Last evening

When it

takes place in

internal parts

these symp do

not come under

our cognizance

Pain is not sufficient to ^{indication} inflam. internal parts

If there is superadded morbid sensibility it

If there is superadded tumefaction & derangement

of function you may be sure that inflam exists

Symptomatic fever is another circumstance

which serves to confirm your opinion

Change in blood itself is an important ^{circumst.} ^{Langs}

in local inflam. = These are the phenomena

the top of blood becomes colorless which constitutes the

buffy coat It also has a cupped appearance

Nov 15th

Causes of Inflammation

Predisposing & Exciting causes

Alterations of temperature most common exciting
cause of local inflam.

Humidity
Suppression of perspiration is a frequent cause
of local inflammation

Heat above 98° F if continued for any length of time
produces inflammation. Poisons taken internally,

Epidemic influences also Morbid poisons -

Inflammation Nov. 15th 1836

Termination of Inflammation
 1st Resolution with or without change of structure
 2 Adhesion
 3 Formation of False membrane
 4 Tropical Effusion
 5 Suppuration
 6 Ulceration

7 Gangrene or mortification
 Resolution It is the object of Physicians to bring about resolution without change of structure
 Adhesion Direct union of two opposite surfaces without the intervention of false membrane
 Nov 16th

Formation of False membrane It takes place most frequently in serous membranes occasionally it takes place in mucous membranes as in Croup & inflammation of bowels

Tropical Effusion It also occurs most freq. in serous membranes

Suppuration It is in some instances a desirable termination pain becomes diminished redness subsides & the tumor points

When it occurs in internal parts pain becomes less it becomes dropping chilliness & weight &c

Pus is a thick yellow fluid without odor & of a marvellous taste

Ulceration Murrate of ammonia coagulates pus but not Apt to take place in mucous textures solution of a continuity of a part attended with some loss of substance

Gangrene It is sometimes the result of high degree of inflammation but not always

pain subsides or altered - dull burning

Inflammation of panes apt to terminate in gangrene

caused by blisters is apt to terminate in gangrene
 Gangrene in young or old persons or in feeble debilitated subjects

Nov. 17th

Inflammation

Gangrene when inflamⁿ of lungs terminates in gangrene
it is indicated by fetor of breath

Constitutional fol. 1st place char of pain becomes
Sym altered & sometimes there is an entire cessation of
pain Countenance changed It is peculiarly pale
great diff of deglutition surface becomes cold
Cold perspiration breaks out singultus subultus
tendinum low muttering delirium Patient
dies comatose

Causes Sometimes owing an epidemic influence of atmosphere
Said in some instances to be produced in
Germany &c by the spotted eye

Modifications of Inflammation

Acute & Chronic inflam.

1st chronic - when it takes place in the viscera it is very
Insidious in its progress Symp. are very obscure
In by far the ^{greater} number of cases it is the result
of acute inflammation

We are the creatures of habit

Force of habit influences the organs &c

Nov. 18th

Subacute inflam. meant to import a particular disease between acute
& chronic inflam. Ex. Bronchitis

Congestion a plethoric state or over distention of the
blood vessels of a part

Treatment Suitable diet & Regimen must constitute
the basis of our treatment

Diet

Antiph. & phlogistic articles of diet
(we merely mean a diet which contains very little
veg decoctions & infusions)

Toast & Barley water gum water & subacid
drinks

Arrow root barley oat meal made

Information Nov 3rd 1836

Blood letting is the most powerful rem in the treat
of acute inflam.

Modifications of the pulse — Frequent pulse
1st year 110 2nd 90 to 100 3rd 90 to 108
3rd to 7th yr it grad comes down to 80
70 to 75 natural frequency of pulse in adult

As gen rule it is more freq in disease than in health
Irregularity of pulse no indication of disease

Rules in feeling of pulse 1st observe force of pulse by compressing it
1st strong & the weak.

1st Strong pulse It exists in healthy system where persons
exercise in open air

2 weak. It exists in delicate persons who take little exer
It is no indication of disease when of natural freq
It is known by degree of compressibility of pulse
When pulse is as high as 120 & you may suppose a
disease exists

Nov 21st wherever

3. Frequent pulse generally speaking it is found in all diseases
when pulse is upwards of 95 it may be called a frequent

4 Slow when not owing to idiosyncrasy it is an indication
of disease
When less than 65 in standing & 60 in horizontal
it is called a slow pulse

Slow pulse is attended with fullness
It is gen owing to derangement of digestive functions
& is mostly relieved by a brisk cathartic & a little abstinence

5 Hard, tense, or bleeding indication of an infam condition
of system There is strongly marked intervals of syst & diast
It gen ranges from 90 to 140

6 Full & soft p when natural in freq is the pulse of health
It is 1 of most favourable indications in the course of
disease

Inflammation

108

Nov. 21st 1836

7 Small or thready it is the very reverse of 6th modification
sensation is that the artery feels constricted
Very apt to occur in inflam of serous membranes
It is always a hard pulse

8. Intermitting or irreg. pulse apt to occur in diseases of heart
Supervention in progress of acute disease - dangerous

Irritable It resembles very much a hard pulse
It is known by looking into previous history of case

Blood-letting It is the most powerful & effective remedy in
treat. of acute inflam
Young children & old persons will not bear the
effects of blood letting as well as middle aged persons
Nov. 22nd Mode of performing the operation of blood-letting
Have all things ready

basin 2 cups 2 bandages 1 1/2 yds. roll of linen
& a glass of water little hartshorn
Bold & decisive blood letting in Acute inflam
should be practiced If faintness is practiced before
we have drawn a sufficient quant place your finger
upon orifice & place the patient in a horizontal
position with slackening the bandage when faintness
has been relieved go on to bleed again

Repetition is to be determined by all the circumstances of the case
If pulse at 10 or 12 remains hard pain continues &
the local inflam is still going on bleed again
Pulse cannot be relied upon but you must take
into consideration all the circumstances of the case

203 of blood is a free bleeding in a good constitution
303 - - - large - - -

Nov. 23rd

Local Blood-letting when there is no
constitutional fever local blood letting is gen
preferable to general blood letting
It has more effect in stopping local inflam
than general blood letting even when carried
to much greater extent

109
Inflammations Nov 23^d 1834

Cupping & leeching compared
We are obliged to use one or the other according
to part from which we are obliged to detract blood
where we can conveniently use cups they are preferable
as they attended with considerable counter irritation

Exorbitant detractions of blood whether spontaneous
or otherwise. It will be fol. by an irritable pulse
& if the bleeding be still persisted in the symps will
be exasperated: Flushed face heat of skin
throbbing & pain in head great throbbing of carotids

1st Case
was a Medical man who received Part injury
from horse falling on him. Had 3 ribs fractured
when reaction took place 16 3/4 blood 12 leeches
to temples when he got home 16 oz more
was taken he was bled again & again
when on 3rd day 12 3/4 blood had been taken
The symps at 8th day was great throbbing of carotids

2 Case In this case the gentleman received
Patient was attacked by pain in orbit to eye
he was a vigorous young man about 27 yrs of age
He was bled freely active cathartic & saline
diaphoretic On 2nd day he was bled again
to amount of 10 3/4 in consequence of pain returning
& on 4th day appeared perfectly well
when he went on some business & pain returned

Nov. 24th Effect of Cathartics in treat of Acute
Local inflammations.

They act by evacuate the bowels, change the secretions
of the intestines & also produce a derivative effect
from Superior to inferior parts of body

Some operate very mild

They are Laxative, ~~gastrointestinal~~ Drastic & others
Drastic Catho. gamboge Jalaterium

Antimonials in treatment of acute local inflam.

Inflammation Nov 24th 1836

Antimonials in treat of acute & local inflammation They are very powerful in subduing it. Among the preparations used the tartar emetic & pulvis antimonialis are preferred.

Tartar emetic may be given in solution in proportion of 1/2 to 3 in table spoonfuls every 4 or 50 min & in powder.

This may be combined with nitre which is easier to digest & its effects in the case of hemorrhages from lungs may be given in proportion of 1/2 to 3 of nitre.

James Powder is a mild antimonial & is used in the treatment of fevers unattended with any very obvious local inflammation.

Blisters - A counter irritant & to be used in common acute inflammation. Blisters should be removed by means of adhesive plaster. They stimulate the organs of organic life said to increase the sensorial powers. I believe that it has very little effect on them. - of Revere

Nov. 25th. Blister if properly prepared should produce its effects in course of from 4 to 6 hours.

Pathological symptoms of inflammation anatomy.

Nov. 28th Inflamm. as it occurs in diff. textures

Rheumatism: Rheumatic fever. Acute & Chronic forms

Acute R. is attended with fever. It is attended generally with rigors & flushes.

attacks the fibrous pulse is gen. small yet it is frequent & hard. In commencement it is attended with wandering pericardium tendons & then it will seize upon some particular part - pericardium, muscles, tendons. In some cases every muscle of voluntary motion. In some cases all the pulse as disease proceed becomes expanded. Blood serum numerous. Perspiration is often copious & acid. Urine is uniformly high coloured attended with a deposit of ceteritious sediment.

111 Rheumatism Nov. 28th 1836

her. is its tendency to shift its situation. Fever generally precedes the fixed pain. It has a disposition to shift from 1 part to another. Free use of leeches combined with warm baths have a great tendency to produce metastasis.

Nov. 29th *Neurodynia* (Rheumatic inflammation of intercostal mus.) You will constantly meet with in practice *Neurodynia*. There is no disposition to come to a crisis. It is most irregular. Sometimes it commences in violent form & suddenly terminates favourably. It rarely terminates in ^{suppuration} ulceration or gangrene. It occasionally terminates in suppuration again in ulcers & not infrequently in caries of bone.

Most common mode is a gelatinous effusion of joint. When once an individual has suffered an attack, predisposition ^{remains} *Causes* ^{causes} Alternations of temperature Moisture Predisposing. Hereditary, by being once attacked a predisposition remains. Mercury increases susceptibility. In these cases it most freq. attacks perivertebra. Occurs most freq. in winter & spring & more freq. in males than females. In plethoric individuals & at the middle periods of life.

Treatment — General principles — It is eminently a phlogistic disease. Nov. 30th Blood-letting is to be employed in the early stage. 16 to 20 3 average quant for 1st bleeding.

Purgatives — *Infusio Senna* with cardamom seeds the this may be laded w. Magnesia. Should not give purgatives in evening. Give *pulvis antimonicus* 5 or 6 gr every 2 or 3 h. & use *pediluvium* in evening. *Spiritus Mindereri* may also be given so as to assist in producing diaphoresis. Warm bath has also been used to produce do.

On 2nd day if symp. not mitigated bleed again. fol. by cathartic. If it has not been exhibited on 1st day. 3rd. Bleeding a 3rd time is as per as allowable.

Antiarthritic — It should be given in large dose = 1 gr. Warm bath in conjunction with blood letting will occasionally cure that disease.

Rheumatism Nov 30th 1836

warm bath has a great tendency to produce metastasis
Metastasis to the heart is the most freq. cause of organic disease
of heart only way in which ~~venous pathology~~ is allowable
is in form of phlebotomy - But use antimony accor.
ding to Italian method

Colchicum Autumnale very much used in Rheumatism
& gout It produces all the symptoms of poison when given
in large doses Wine of seeds of seeds is the best
official Prep.

Gallate of Veratrum $\frac{1}{4}$ to $\frac{1}{2}$ gr. eq. to 20 or 30 qtt of wine of seeds
Not give it in early stage but reduce system by bloodletting
Antimonials for 2 or 3 days

Formula { Water 13
Sulphur Mag. 23
Carbonate - 23
Wine of seeds 20 or 30 qtt } repeated ev. for
6 hours

Opium - Most freq. used in form of Dover's powder
or in form of solid opium in combination with calomel
occasionally it produces good effects in early stage but
much more freq it does harm and 1st & 2nd day
It will sometimes produce a quiet sleep when given in 4 or 5 gr doses
in combination with calomel

Cinchona officinalis I have used it frequently and occasionally
with advantage but more frequently fails It may be
used where other remedies have failed in seeing it

Local Rem such as blisters and other counter irritations are
not benefited in 1st periods of acute Rheumatism
until inflam symp have been subdued I shall
therefore go on the spears of

Chronic Rheumatism

Distinctive character between acute & chronic form
is that former is attended with fever latter form is not
It is mostly seated in some particular part It mostly the
sequel of acute form though not always

Remedies diff. materially in 2 forms Gen. bleeding seldom
required Local blood letting is to be preferred

Rheumatism Dec 1st 1836

Stimulating drugs very much used in most of the Cures
of Chronic Rheumatism is the *Aprobrilum Medicorum* of
medical science crown of glory to quacks & old women

Gum Guaiacum sustains the highest reputation & should be
giv in Aromatic ammoniated tincture Giv in dose of teaspoon
full increased to table spoonful 2 or 3 times a day
It has the effect of keeping bowels open

Plum Jer. It is the best remedy that can be employed when giv in
large doses

Formula } ol. Jer. 15
 } Honey 43 } table spoon. ev 2 hours
 } Grad add water 83

A convenient form is to rub up 4 oz of oil with yolk of egg
& grad add ~~83 water~~

Balsam Copiva has very nearly the same effects as ol. Terbinthina
When it has been produced by exposure after mercury has been used
confine your patient to bed and give large dose of Compound
decodation of Sarsaparilla

Carbonate of Iron in dose of 3 to 3 has been strongly recom
Mercury so as to produce gentle ptyalism has a very good effect
External Remedies are very numerous

Both in Sciatica & lumbago treated by counterirritation as blisters

In France they use a more powerful remedy viz Moxa
formed of cotton dipped in nitrate of potash I prefer the Spunk

Sinapisms are very useful remedies

Cantharides boiled in turpentine is a very powerful refrigeratory rem
Burgundy pitch plaster very com remedy

Dec 2nd Friction with hand is a very good remedy

Sulphurous vapour bath It brings on copious perspiration

& makes the skin very red in fact it acts as a rubefacient

Acupuncture In some cases it acts almost like magic

Gout December 2nd 1830

Distinction between Gout & Rheum

They are both inflammations & both attack the same structures
 They both change place &c
 Rheum produced by cold & moisture Gout is not but
 is more frequently transmitted from parents to offspring Gout
 is dependent on certain modes of living
 It is preceded by signs of indigestion Often an attack
 of gout attacks health it impairs Gout so in rheumatism
 A rheumatism attacks the larger joints attacks all persons
 young and old alike

Dec 5th Gout is emphatically a disease of the rich and wealthy
 They are both attended with severe pain
 Diff. 1st Signs Situation Rheum in large joints in
 small joints. Ages. Terminations.
 It occurs in robust plethoric persons Most common
 cause is high seasoned & animal food
 Habitual drinkers of alcohol do not have the gout
 The drinking of wine with high living have a tendency
 Exciting C. depressed

Comes on in this way There is symp of indigestion
 the foot feels distended This is attributed to tight shoe
 he goes to bed & is waked up in middle of night
 with great pain in foot which keeps him awake
 On exam. there is a pulse and glossy red appears
 On the 3rd night the pain is agonizing The toe
 cannot be touched with causing pain By this
 time very much distended red & great pain

First fit seldom continues more than 3 or 4 days before
 the symptoms begin to subside & will not return
 if patient is cautious under a period of 12 months

The whole of texture of joint suffers in this disease
 Disease is primarily seated in digestive organs

It is an hereditary or acquired disease

Predis. C. They are frequently eating & drinking
 Great Profligate treat By a proper mode of diet & regimen
 to may prevent an attack of Gout where a predisposition
 Paroxysm choice of 3 modes 1st to let it run its course 2nd to cut it short
 n 3rd to adopt an intermediate course 1st purgatives patient's plan
 n 2nd to use cold applications to cut it short Another new to cut it short
 n 3rd modicenal

Simple Inflammation Dec 6th 1836

Dec 7th Inflammation as it occurs in the Brain
 Ignorant of the ultimate structure of the brain
 We can trace no connection between their mechanical forms
 & their functions Brain is the material organ of thought
 It is the source of thought & sensation
 Inflammation gen extends from ~~one~~ texture to another more
 than in any other inflammation
 Pain in head is no evidence of inflammation of brain - nor
 is derangement in functions of brain Even the pulse
 is very apt to depart us

Dec 8th It is impossible to say whether this or that mens. is inflamed
 20,000 annually die of inflam. of brain in this country

Cerebritis or inflam. of brain and its meninges

1st Modification runs its course
 Threnitis Idiop. is a very uncommon disease It comes on
 very soon in young by exposure to sun & cold
 substances applied to skin Rubric large glands
 of electric drinks at a time & so of the above the person
 Repulsion of cutaneous disease

Symptoms vary much according to circumstances by which it was
 When it arises by very slight cause
 1st group fever That is person will be rather I
 The person will say head feels as if a band of iron was bound round the
 The eye has a peculiar appearance

Hydrocephalus or Meningitis it occurs in children It occurs
 most frequently during dentition They are common

Dec 9th the most common of the diseases you will be called upon to treat
 Chronic Hydrocephalus

Next modifications of which I shall speak
 is when it fol. suppuration from Cea
 It is 1 of the most common sources of this disease
 next form

Brain softening of Brain
 Dr Wilson post directed the attention of the profession to it
 Most of cases of ~~severe~~ apoplexy & hydrocephalus & convulsions
 are owing to it

Cerebritis Dec 9th 1836

It consists of softening of brain forming a soft putty like mass when you pass hands of scalp over ~~softened~~ ^{softened} brain it adheres to it. It will not in healthy brain. It will sometimes present other appearances as false meninges in ventricles, formation of matter & fist in brain &c. In some cases there is no doubt that inflammation exists in the brain. Occasionally softening is the consequence of inflammation. At other times there is no sign of inflammation. Like gangrene or mortification it may be the result of inflammation or may be the result of stasis in the circulation. It never occurs beyond 30 yrs of Age. Prognosis. Another modification is disease of brain connected with tubercles occurring in its structure.

Dec 12th 1836 Softening of brain most freq takes place in corpus striatum.

Prognosis

Cerebritis in all its forms is highly dangerous. Prognosis says that he never cured but 2 cases of softening of brain. Great diff is the diagnosis. Rule. When we can ascertain that inflammation exists use active treatment.

Diagnosis

Before disorganization has taken place. 1st violent pain in head. Superadded Delirium in early part of disease. They are not direct proofs but they are strong signs. Paralysis. Appearance of stools. In hydrocephalus they are green & resemble parsley.

Dec 14 Coma & convulsions taken in conjunction with the other circumstances are the proof that inflammation of the brain exists.

Treat it is one of the most inflamed diseases & req the most powerful antiphlogistic rem.

Blood letting must be carried to an extent that would scarcely be allowable in any other disease. It is the heroic. In course of 48 Child of 15 or 12 yrs 50 or 60 oz. Where veins are small you may bleed from jugular vein. When gen blood letting has been carried to sufficient extent Local blood

117

Rever's

Notes taken from Dr. ~~Rever's~~ Lectures

Nov 5th

on Practise of Medicine

~~Human anatomy is a science which is the foundation of the~~
~~science of medicine~~

Treatment of Cerebritis

Cathartics come in next to blood-letting
Where stomach is irritable calomel is the best - give
in large doses & when stomach is composed if bowels
have not been evacuated use some active cathartic

Cold Applications to head is a powerful & good remedy
By means of bladder or more powerful is by means of wet cloth
As far the most powerful is by what is called the cold cloth
over the head. It will reduce the furious delirium that attends

Blisters have been considered among the doubtful rem

In some cases they are injurious. After the disease has been
treated actively & disease appears at a stand by shaving scalp & applying
blister to scalp I have seen the most beneficial effect -
occasionally it is the reverse. If it causes irritation

apply simple cerate. Sprinkle with acetate & sulphate of morphia
Mercury It is by no means so good a rem as was formerly supposed
Its best effect is to evacuate the bowels

Digitalis Purpurea In this disease I have never seen it produce
Fourth rem in power is Gartrate of Antimony

The rem are few energetic & obvious. An early & unfavourable
Prognosis has very unfavourable effect upon patient & also
upon the Practitioner. It is never too late to despair

Rachialgitis Dec^r 14th 1836

Dec^r 14th These disease connected with inflammation of vertebral
column

Medulla is to be considered a part of brain protected by same men
The chord is liable to same affections as brain
Chronic circumscribed inflammations are liable to take place.
Many nervous diseases depends upon this chronic inflammation

1 Inflan of spinal cord or membranes known by symptoms

2nd Where disease appears to exist in some other part

3rd Next I shall speak of the various nervous diseases attributed to this source

Abstract Symp. 1st Where the symp obviously indicate the seat to be in spinal cord

severe pain in the track of spine
pain sometimes remains fixed in some part of spine

Pain sometimes begins in cervical & extends down

spine with numbness of thighs diff in voiding urine

Where it is in cervical portion is affected you have pain

shooting up into head Stiffness of tongue Head fixed

Where dorsal portion is affected you will have dyspnoea as pain

in epigastrium under left mammae symp of dyspepsia

Dec 16th After death the true source is found to exist in medulla

Cerebrum & cerebellum seldom involved in conjunction

more acute form Rigors, flushes,

Arise from exposure to cold occurs most frequently in

persons who remain in bent position Scrofulous constitution

They are indicated by their sensations & confirmed by exam after death

Neurosis of Culler Dependent upon disease of brain & medulla spinalis

2nd class Those in which the patient gets well & therefore the disease is inferred from

morbid sensibility & derangement of function

How do you know the existence of morbid sensibility in this part

Answer & Prepare Pain under left mammae sometimes under right mammae

Dec 17th Monday Great emaciation; freq palpitation of heart supposed to depend upon disease of heart Upon examination you will find tenderness to pressure in left axilla or 1st dorsal vertebra

Case of Taylor health became impaired weight about 140 lbs. strength became so weak that he was obliged to quit his business on examination the ~~case~~ compound nerve was found to give rise to chorea chorea of face & upper limbs was treated with a blister applied over the heart the young man speedily recovered

Notes Taken from Dr. Revers's Case

Case 1st - A lady, one the fourth evening after being confined of 4th child - through the course of the night on hearing one of ~~the~~ her children crying in a distant room in the house got up out the bed in order to go to attend to the child, the entry leading to the room through which she passed being covered with a glazed carpet & the lady after having reached the room door considering it a very imprudent act in her immediately returned to her bed - She was immediately seized with a violent chill. Likewise followed with severe pains under the mammae head & lumbar regions - with a suppression of the milk & lochia - Medical attention of the highest rank paid to her - she had under gone the every possible mode of treatment - and at last had come to the conclusion never to consult a medical gentleman again - She at last resorted to her own mode of treatment which was as follows - when the lady felt those violent returning she would wrap her self up in wollen blankets & apply warm bricks to her feet - on happening to be when this lady was & hearing her express those violent pains she had been subjected to for 10 or 12 years past I requested her to let me examine her spine by pressing upon each vertebral process I ascribed it to be a disease of the cornu poma nervi - which I then cupped her on one side of the spine and applied a blister immediately over the vertebrae which I found gave her great sensation of pain - the lady in a short time began to recover & never has experienced those severe pains since -

Case of - who was affected with the paralysis of the left arm also with violent pains in brain which was found to arise from those nerves - Cupping & Blisters - was resort to which relieved the man in 3 weeks though he had laboured under those pains 2 years with

John A. Morris

1836 Derangements in the functions of the Rectum

Dec. 20th Views of Dr. O'Brien. Does contain in coecum
 & colon there is inclined to exist a stricture at termination
 of sigmoid flexure of colon in rectum. This stricture
 sometimes prevents injections from passing up.
 There is a pouch in upper part of rectum sufficient to contain
 only 1 or 2 fl oz.

In affection of Rectum bladder & uterus. The sympathetic
 between these organs are to be taken into consideration.
 A longer tube should be occasionally employed.

Dec. 21st Neurosis

Hysteria. It generally occurs in paroxysms.
 Person appears to be in ordinary health. From some
 sudden cause, she will become dull gloomy & not sleep
 & then she will burst out in a fit of laughter. These
 will alternate for a while, when she will be thrown
 into violent convulsions & voluntary violence.
 These attacks are sometimes slight & sometimes serious.
 It occurs in persons of all constitutions. Quite as freq.
 in robust as weak. Very freq. attended with suppression
 of menses. Catamenia are in some way or other
 deranged. There is preceding or commencing with it pain
 in lower left mamma. Pains in loins. Palpitation
 of heart sometimes preceding sometimes accompanies the paroxysm.

Treatment

As ipecacuanha has rather the effect of increasing the paroxysm.
 You must prescribe according to symp. If they
 indicate bleeding resort to it. If not req. Antispasmodics
 will have the effect of relieving the paroxysm.
 Rule - need not not to be excited when called to a case.
 Chorea Sanativitis. It consists of convulsive
 twitching of voluntary muscles. It is connected with
 some disease of spinal medulla.

Pneumonia of Respiratory Organs

Laryngitis 1st as it occurs in childhood & 2nd as it occurs in adults

1st Laryngitis of childhood a Croup more fatal in childhood than in adults. It occurs at all seasons all climates most frequent in winter & spring.

Symptoms

After a damp day it is preceded with a cold it is just to bed and wakes up with croup. At other times child goes to bed well & wakes up with symp of croup.

Children soon recover but have had a great many comparisons.

It is remarkable for coming on after short inspiration.

It is between a cough & a grunt. Perhaps the child will be spirited & playful. Ev few minutes it will have this cough. At this period it is quite a local disease & is very easily cured at this period. It is only attended with symp of common catarrh.

the croupy cough begins hoarse voice.

the fever attacks the patient - attended with frequent hard pulse. Feverish symp become violent & incontinence.

becomes suffused & swollen but is no longer of a bright hue. extremities grow cold. The suffusion becomes intense.

The child is seized with convulsions & dies. Does not form 2nd to 5th day & sometimes within 24 hours.

Sometimes inflam great extends down bronchia child has not great dyspnea in drawing air in throat & much

inflammation in throat.

Inflam of larynx & trachea & bronchia & false membrane & very often formed.

Pathology - 3 stages

1st Forming stage from commencement to the period when symptoms begin to take place.

2nd stage when symp begin to take place.

3rd stage when the lungs begin to fail in their functions.

There is a marked secretion which is seldom met with & violent effort to throw it off & into diff in adult.

Croup

Dec 1 1842

Great well known diff in three different stages

1st period marked with danger of attending to
2nd stage

3rd It is most formidable in this period

Dec 29th Treatment of croup

It is rapid in its attack rapid in its course & affects
parts important to life

Should attend to every hoarse cough in children

If it occurs at midnight should call Physician immediately

Treat it with great certainty at 1st period by means of emetic

Tartar emetic is decidedly the best emetic dissolved in water

child 6 or 8 yrs old 1 table spoonful of solution proportion of
4 grs to 3 The dose would 2 grsShould not be repeated under 50 minutes It is to
its powerful antiphlogistic effects that we may attribute an impor-
tant benefit derived from the remedy

If the antimony does not evacuate the bowels in course of hour

1 or 2 h after vomiting give a purgative & Calomel is

the best remedy that can be used. It is not unpleasant

to take & it alters the secretions

Keep the system

slightly nauseated for at least 24 hours

2nd stage Emetic not the 1st remedy in this stage

Most powerful antiphlogistic must be used

Blood letting in some form or other ~~very~~ very young children

have to bleed from external jugular Some times best to bleed

by means of leech to top of sternum 3, 6 or 8 3 from child

3, 4 or 5 yrs of age There is here a race between the disease

& the remedy employed - they should be powerful

3rd stage - Consequences of bloodletting is sometimes immediately
fatal Tartar Emetic again becomes the main remedy

Blister use them though I have seen very little benefit

We are in the habit of use leeches in young children when

we bleed letting

Polygla Senega not found useful in my hands

Calomel improper only as a purgative

has been a great remedy in its treatment

3rd stage Emetic warm bath

Laryngitis of Adults December 30th 1836

Identically the same disease with Croup of children
 Washington fell from this disease
 It usually begins above the glottis constituting subglottic tonsillitis
 attended with pain & swelling in throat & diff. of swallowing
 It grad extends to Larynx attended with a harsh shrill
 ringing cough

Case Gen. proves fatal from the 1st to 3rd day
 Gen induced like Croup by change of temperature

Case of Gen 17 Decr 1836

Croup appears to be a mild ^{disease} when compared with Laryngitis of Adults
 Great Gen Bloodletting ^{instituted} and other decided antiphlogistic remedies
 Fatal effect is produced by swelling about larynx stopping passage
 of air to lungs Laryngotomy Bronchotomy

Seldom performed in children More likely to prove successful
 in adults In Children the larynx & ramifications of bronchiae
 are blocked up with false membrane In adult
 the inflame is extremely limited seldom extending below the larynx
 There is seldom a deposition of false membrane
 If fatal symp are advancing after prop treat. perform the operation
 Relief not being obtained is generally owing to the operation being
 put off too long

January 2nd 1837 Inflammation of the Pulmonic
 organs

Diff modes of Examining the Chest

Hitherto only 2 modes viz the touch & the sight
 The sense of hearing has been very little employed

The chest is the most sensitive of any part of the body
 Percussion of the Chest 1844 used by Steinhilber

Resonance of chest diff in diff parts Use all the
 fingers to strike the chest the finger of the other hand should
 intervene Before striking direct to fill his lungs & then while
 the lung is in diff states When the anterior part of chest
 is examined shoulders should be elevated & drawn back
 where there is no resonance in a part mark it in thereby
 & also compare it with the side

Examinations of the Chest, Jan^y 1837

1837 vainly presumption & the very spirit of quackery
 January 3rd Mediate Auscultation & Percussion
 Percussion introduced in 1763 by Auenbrugger

It is made under 3 diff. forms 1st when lungs are distended 2nd when they are empty & 3rd when they are in mid state between distention & contraction

Immediate Auscultations attended with certain difficulties to remedy which the stethoscope has been used to remove them - Mediate Auscultation sounds by voice should be examined by making tube of continuous size It is best to have but 1 fold of linen superimposed which should be made smooth 3 modes

1st in relation voice 2nd Respiration 3rd in relation to heart
 1 Examination of voice

2 Explorations of Respiration It should be used in form of trumpet Should succeed to percussion

Sounds are generally louder above the clavicles Much more distinct in child than adult & from that circumstance called puerile It exists sometimes in adult & is freq the consequence of certain portions of lung being impermeable to air

When we find no respiratory murmur in certain portions of lungs it is owing to some obstruction to passage of air When we find respiratory murmur nearly natural all over the chest the disease is not dangerous even though the signs are violent

On Regard to Circulation

An appreciable difference exists in regard to time between systole of heart & pulse at wrist

January 11th

Bronchitis Inflammation of mucous membrane of lungs Catarrh & bronchitis only diff. in degree It is a violent disease & sometimes terminates suddenly

Bronchitis Jan 4th 1837

Symptoms Symp. fever sore throat running at nose
cough coming on in paroxysms Tightness about the chest
Change of temperature apt to bring on a paroxysm
of cough. Mucous has the form of the air cells

Character - Pain in head is very characteristic
When he coughs or stoops he applies his hands to head
Tongue is gon large & hanging over the teeth
There are reg. exacerbations every day worse at 10 o'clock P.M.
Expectations of sputum & mucus mucus of lungs have a great sympathy
existing between them

Character Head ache increased by jarring cough the
Tightness of chest is very remarkable
Instead of being active it is weak & unimprovable for it
blood letting very well borne Another is the copious
expectoration Last circumstance is that the disease
soon breaks down the energies of the system

January 5th 1837

Treatment

At early period the most decided antiphlogistic means
must be used Patient, cannot bear blood letting to great
extent Should make a small incision & to a moderate
extent to gether with cups & repetition of the remedy
Emetics They are very important rem & is indispensable
in children **Antimony** Sometimes quills

Purgatives & opacac is pref in children
The milder of this remedies so as merely
to remove the peccant &c

Among the remedies is Calomel usually give
in divided doses 2 or 3 gr Sometimes 1/2 gr to produce

Blisters should not be used in early stage but are advantageous
at a later period Small blisters & frequently applied

Opium has the effect to produce constipation & stimulate
the system After 3 or 4 day a dose of opium viz going
to bed

Diet regimen patient should drink freely
of mucilages barley water gum water plover
seed tea &c I shall speak of diet under head of chronic

Chronic Bronchitis

It comes on gradually / It is only attended with cough in morning

One of the effects of this chronic cough is the enlargement of bronchial passages

Acute Bronchitis is exceedingly apt to degenerate into Chronic form. Patient wakes up in morning when cough begins & continues sometimes until the patient becomes perfectly exhausted.

Slightest change of temperature will bring on a chill even in getting to bed.

The symptoms grow increasing which may be noticed from week to week. Digestion becomes disordered

bowels irregular. Surface pale. Pulse becomes affected about 100 in horizontal posture increased 10 or 12 beats on rising up.

Patient becomes worse at certain periods of day. Hands & feet burn. Cheeks flushed &c.

There is an exultation of intellect which even the patient becomes sensible. Some organic lesion is going on. I have seen 2 peroxysms

of hectic in course of day. There is night sweat with cold hands & feet. They sometimes expectorate

a qt. of green mucus in course of 24 hours. Lecture Review

Chronic cough. Copious expectoration. I am told of emaciation & finally hectic fever.

Stomach cutters glass cutters & are particularly liable to this disease. Healing hemp hat skins hat chelms hat chelms

Hemoptysis is a dangerous symptom. 3 modification of this disease

progress. Gen. syst. of indigestion for months. dyspepsia, pains in side slight cough.

Sallow skin palpitation of heart and various symptoms. Many of these cases are complicated with spinal disease.

Both ~~Wormen~~ Lungs do not collapse. Principal marks of disease are to be found in bronchus

they are loaded with mucus & very much dilated. Pleura & Pleura Pulmonalis are adherent.

Treatment. Gen blood letting is it proper? as we are in habit of seeing it. It is not warrantable

where necessary. Local is to be preferred.

Bleeding from arm is necessary under particular circumstances.

127

Chronic Bronchitis Jan 6th 1837

Counter Irritation among the most valuable remedies
Tartar E. ointment, caustic seton may answer a better purpose
Where the pleura is inflamed a blister is the best
Emetics are among the most common & occasionally useful
Stomach is with diff. associated
Mustard Specific
Mustard for - salt & water
5 gr mustard
5 gr pulvis Specific is the
one I mostly use in children

Expectorants
Vinous tinct of Colchicum acts on bowels It deserves that
deserves more notice &c
Balsams 1st Balsam Capivi never seen any good
result from it objectionable
Anell done of eol combined with Opium & one of the expectorants
is the best remedy that I have ever used

Formed { Op 5 to 12 gr
Squib 5
Emetic 1
Cal 4 gr
made into 8 pills
Take 1 at night & if the case is troublesome give
another in morning

Where case is attended with copious expectoration
give 4 quinine with 4 or 5 grs laudanum
Sometimes they bear it best in form of emulsion with starch
& laudanum

Pirra or gapes in detail they have been laid aside
vapour of tur is only one that is attended with any advantage
Night Hæmorrhage

Elix V made like lemonade
& let the patient drink freely
Diet & Regimen is very important
Farrinae & milk diet is to be preferred avoiding all
stimulating articles In advanced stage you are obliged
to leave his own whims if you find that they disagree
Prohibit them or allow them only in moderation

Chronic Bronchitis Jan 6th 1837

Regimen on the last exposure the patient will have a chill
Rooms should be kept of reg temperature by an open fire
place Riding in carriage has a heating effect to

Jan. 9th 1837 Gen. speaking Local bloodletting is to be super
mild emetic Salt water 5 to 6 gr ipecac
rubbed up with mustard seed Cathartics
merely keep the bowels open Balsam Cap. not useful
Sulphate of quinine with Ladannum will reduce the expectoration
It will sometimes stop it to much when given in enemas
Regimen & diet Great dependence is to be placed upon
them Temperature has a great effect Mr. Jones Store
The symp are such that we can lead to a great certainty
the texture

Inflammation of Parenchyma of Lungs.

Pneumonitis Jan 9th 1837

smallest symp fever pain in the chest
pain in chest is deep seated circumscribed & is in a much
more increased in it than in Bronchitis when patient is
placed horizontally Patient is obliged to have head
shoulders elevated Respiration is diff with sensation
of cord around throat Cough is not prolonged but
is merely a short grunt Face pale profuse
sweats about upper part of chest without relief
Expectoration at 1st small about 2 or 3 d day the mucus
is intimately mixed up with blood called purpurine
expectoration - This is the Pathognomonic sign generally
Sometimes it is absent In more advanced stage
it becomes yellow streaked with blood
pulse is full round & hard Char of tongue in the
flaming large & relaxed is of a coriaceous pink
great thirst & inappetency Fever increase at night

Assaultation & preparation is

Pain & unperf. resonance in same part would enable to know the extent. Stetho. Exp. is a very valuable means. Air passages being diminished the resonance is lost. Respiratory murmur is attended with a crackling sound - from air passing through mucus = crepitating rales. In advanced stages

the parts become solid & of simpler lives. They terminate in
reparation ^{decomposition} & inspiration & organization

Hyperemia or engorgement of the lungs

^{or} Hepatization. Sometimes of colour of liver changes from that to a yellow
It is next in frequency to Hyperemia

Sometimes put it fine in filter into substance of Cuz.

Abies very rare

Gangrene necessarily around

to be returned in care of 1840

Adonia Lungs have a dark greyish appearance & when cut water issues out. Lacunae ^{scarcely} apparent at latter period

Occurs most frequently between 45 & 50 years

Upreg. in males than females. Most fec. in winter & spring.

prefigured me, and the exanthematic. The attack
pre-disposes to another it has occurred 28 times in one individual.

Treatments Blood-letting all agree that it is useful

It should be used very freely

Can. 50 the very important July 2^d of the 2

Restrictive circumstances,

Char of Pneum. is fixed opaque & deep ^{red} coloration from
since ~~change of structure~~

gives
Hepaticization. Change of color
red & yellow

ned is the first step after curing our
light coloured hepatization is process of nature to restore
the lung to its natural state.

In bronchitis disorganization is slow while in pneumonia it is exceedingly rapid

Treat. It is a decidedly phlogistic disease. It is proper to deplete the system freely by bloodletting which is the main remedy. If symp. keep up after 1st bleeding bleed again freely & fairly & continue to use the remedy for 3 or 4 days if the disease is not subdued.

Pneumonitis January 18th

Tartrate of Antimony - 1 gr ev 2h - increased to 2 or 3 grs
 I think that $\frac{1}{2}$ gr to 1 gr ev. 3 or 4 h is quite sufficient
 Inhaling warm water has a good effect It promotes the
 process of expectoration

Counter Irritation Vessication should be used
Calomel 3 to 5 gr repeated every 3 or 4 h
 Can very often cure the disease by this remedy with as
 great certainty as by tartar emetic **Pneumonia Billroth**

Chronic Pneumonitis is a very rare disease
 Hepatization is sometimes attended with tubercles

Acute Pleuritis

Acute pain in some part & appears to dart from fore
 front at lower part of chest also Pneumonitis
 sensation of burning over surface of lungs Cough speaking
 or sneezing increases the pain

Position he places in some recumbent posture & confines
 himself to that posture In Pneumonitis he lies upon
 Back In Pleuritis he lies on opposite side or on back

The reason is that there is morbid sensibility in the part
 In Pneumonitis the lungs are infl. & increased therefore he lies
 upon affected side Pleuritis is hurried he is
 unable to speak abdominal & panting respiration
 Pulse is strong Countenance is full suffused; when the
 inflammation extends to pleura lung diaphragm & synch worse
 Expectoration scanty

Character Lancinating pain of colourless expectoration
 Pneumonitis fixed deep seated dull pain on opposite side in early stage
 Describes affected side or back Prime juice expectoration scanty 3 days
 Progress variable sometimes fatal in period of from 2 1/2 to 3 days
 mostly continues 2 1/2 days

Cause one attack does not dispose to a 2nd
Post Mortem red points over surface of inflamed pleura
 are not thickened though has that appearance but is owing to the points
 & to formation of false membrane

131
Pewrites Jan. 10. the 1837

Dropsical effusion instead of being the last of the series of phenomena
 Jan. 11th 1834 occurs at onset of disease. In some cases it is white transparent
 or yellow. Whey like &c. The fluid uniformly that
 the serous fluid is increased. Formations of plates & membranes
 The attack protects in some degree the person from a 2nd
 attack. When the fluid is small - quant it occupies
 lower part of chest & occasional little inconvenience.
 It sometimes crowds the lung into space not larger than fist
 Qu is sometimes effused instead of dropsy = Empyema
 In this case the affected side is enlarged more than in
 case of dropsical effusion
 Adhesions are among the most common appearances in effusion

Adhesions are among the most common appearances in diseased
 credit. Highly phlogistic and like Pneumonia it must
 be treated with decided antiphlogistics. ~~Adhesions~~
 The principles laid down for the treatment of Pneumonia
 will suffice for that of Pleuritis.

Jan 2th Chronic ^{will suffice for Heat of Pleuritis} ~~Pneumonia~~ Pleuritis
 Remarkable Character Contractions of chest

Remarkable Characters Contractions of chest

Pneumothorax = effusion of water a pus & and air
the remarkable char when water is effused is ρ_{50} phony

Dr. Whist diff from the influm I have already described
It is the only one neplary palat. Lot mortem you
find tubercles in the lungs instead of hepatization dropsy

Jan. 13th 1837 Hydatids are animals of the very lowest degree of vitality. Tubercles are originally formed

from small glands in lungs from respiration
that class of chloro C catarrhes with consolidation
& after death tubercles in lungs

Respiration is very frequent. Patient can take a full breath but can not retain it without a heaving cough. If superadded there is delicate conjugation emaciation little muscular energy hectic fever — you may be sure that Phthisis Pulmonalis. Recurrence of hemoptysis is a strong diagnostic of Phthisis.

132 Phthisis Jan 16th 1837

Tubercles in some way or other produced by inflammation
 A smothering disease begins abruptly Hemoptysis may be the
 1st sym

Mixed & 1st process of digestion
 are the only functions that remain perfect to the last

Low delirium occasionally supervenes just before death but
 more frequently he is calm & dies quite unexpectedly

Diarrhoea takes place towards its close Occasionally he
 will appear for months together to be convalescent

Gen Char 1st as to evolution of tubercles disease

is Phthisis unless tubercles exist Origin always
 originates from exposure to cold Duration 3 or 4 weeks

to as many years Cough is a variable though one

of the most predominant symptoms Most troublesome

at night It is sometimes very troublesome

Expectoration early scanty transparent viscid Late

it is thick, yellow copious sometimes mixed with blood

Hemoptysis comes on frequently as the very 1st symptom

Fleeting fever is one of the most frequent symptoms

at 1st only one paroxysm in 24h & that in evening

at advanced period 2 paroxysms occur in day

Diarrhoea After a violent paroxysm the patient

will sometimes experience pain in side & tremor become cold

& the patient dies in course of 24 hours

To distinguish it from Chronic Bronch

Char of affection &c cannot be depended upon

one case - Dry hacking cough Muscular debility & after

a little while the expectoration becomes purulent - is sufficient

to indicate that the disease is tubercular - May be

deceived even in this case as when abscess forms in liver

Galloping Con & consumption is tubercular disease where

there is a great many tubercles developed at once

Diagnosis is important If satisfied that the case

is Phthisis we can do nothing but smooth the patient's way

to the grave Patient should not be sent abroad

In doubtful cases the stethoscope is a very valuable instrument

overrun with a rattle indicates a cavity in forming

Lectology is owing to excavations dependent upon tuberculation

the voice reverberates in cavity

Y. *Phthisis* Jan. 17th. 1837

It is impossible to distinguish between Chronic bronchitis & Phthisis without the aid of percussion & Auscultation
Mellencosis or Laryngeal Phthisis

Hoarse cough change in voice & great diff. of expectoration
owing to muscles of larynx being affected
hence invariably connected with it Phthisis

Causes of Phthisis Predisposing hereditary

sometimes acquired from irreg habits in diet clothing
Pneumonia has very little effect in producing it as it occurs
most freq in lower part of lungs while Phthis occurs ^{part of lung}
freq attributed to catarrhs or common colds. But I think
it has very little to do in causing Phthisis

It is most freq transmitted from parents to offspring

Another Cause is the depressing passions

Jan 18th 3 one or 2 other causes Climate has the effect
of producing them & when they are developed of increasing
them Age Most com period of development is between
16 & 30 yrs It is more likely to be developed in cities
than country Small Pox & measles particularly
latter is liable to produce the disease

Preguidance abroad is that the disease is contagious. It is not Contag

Impure atmosphere dust &c produce the disease
Do tubercles ever absorbed

When small number of tubercles are formed they may come to maturity
be expectorated & the patient get well

Meat

Wherever you hear of a great number of remedies being used
we may by good management stop or at least check the disease
Bleeding in Phthisis no well informed practitioner will
use it except when it is connected with inflammation

When hemoptysis occurs it does require bleeding with
other remedies as Nitrate of Potash Dose 15 to 30 grs

Counter irritation Canthar, Moxa

Where there force & deep seated pains - permanent irritation & Emetics

Where stich occurs a small blister
Digitalis Purpurea or Foxglove

1837

134

Sharpsbury N.J. Aug

Pneumonia Pulmonalis Jan 18th

Mercurial preparations useless in scrofulous disease
 Prussic acid In small doses increases digestion & patient
 is pleased with it at 1st. But after it proves useless
 Residence in a low marshy situation Just like all
 other remedies it turns out to be useless
 All the gapes have had a momentary reputation useless
 Sea Air = Laennec Iodine

Fever

Jan. 19th (Thursday) 1837

Some attribute all fevers to some local inflammation.
 Def. of Fever - Sharpy Rigors, increased heat & increase in freq
 of pulse. These sym occur in almost all idiopathic fevers
 Idiopathic Fe Where there is derangement in all
 the functions of the body without out any obvious local inflam.
 They are also char by rigors increased heat & freq of pulse
 In many cases of idiopathic fever
 When the fever is attended with obvious symp of inflame or
 of brain &c it is called Phlegmemic
 Endemic = goitre Intermittent Remittent
 Epidemic = those diseases which occur at intervals Catter
 they are referable to certain conditions of the atmosphere
 Sporadic = reverse to epidemics one or two cases in place
 Contagious = diseases which produce the same diseases
 in healthy persons.
 Infectious = disease which arise from crowding together
 a great number of persons & which when produced
 produces a disease similar to itself

Predisposing cause of Intermittent &

Remittent Fever

During process of decomposition of animal ^{& vegetable} substances
 Malaria - It produces intermittent fever
 Presence of organized being in a state of decomposition
 certain degree of heat & moisture

100 F moist pavour the decomposition of animal substances
 110 or 120 F most pervious to vegetable substances

Intermittent & Remittent Fever

January 20th 1837 I. & R. are never found where the process of decomposition is not going on. Where you have low land marshy situations you are very sure to have Inter-
Predisposing cause of intermittent & remittent fever is process of decomposition of animal & vegetable substances

Jan 21st

Nature & properties & C of Predisposing cause of I & R fever = malaria Produces its effect in period of from 20 & 3 hours to a month or two

Those who have been exposed to it for a length of time are not likely to have the disease produced.

Youth vigorous constitutions exercise in open air appears to have a tendency to predispose to its effects
Malaria produces inflammation of abdominal viscera & various other diseases

Jan 23rd 1837 Simple Intermittent Fever

1 striking char. is their periodicity They recur with great regularity At the malignant form they are not regular

Causes — Only predisposing cause of I & R is malaria
Worms suppurations of catarrhs & C may act as occasional exciting causes after person has been exposed to the predisposing cause = malaria

January 27th 1837

Remittent Fever

It is divided into simple & malignant forms
Symptoms resemble those of intermittent fever
Sometimes come on abruptly

Cold fit is gen. shorter than in intermittent

Hot stage is more marked longer

Quin in heat comes on early in this disease

Delirium sometimes occurs at the very outset of the disease

136

Remittent Fever Jan 27th 1837

Patient becomes taciturn, nervous &c occasionally it is the reverse Gastric uneasiness intense burning thirst nausea vomiting of bilious matter = It does not mitigate the headache

Headache & vomiting come on about same times In the worst cases there is no evidence of bile in the secretions The urine small in quantity & acid In worst forms there is entire suppression

Respiratory functions affected &c In cold stage pulse is thready & small Tongue in progress of case the tongue becomes yellow edged red &c

One of the most uniform symp is thirst The more he drinks the more the thirst seems to be expressed Bowels are very costive Moved with diff Fumica albuginea yellow In some cases the body becomes jaundiced

Gen. char. are the same as those of intermittent They gen terminate favourably ^{in 3rd or 4th day} on the 3rd or 4th or 5th day Intermittent Simple Remittent & Bilious Malignant Remittent or yellow fever These I think all produced by malarie

Continued Fever February 3rd 1837

They gen occur in cold weather commences in fall & continue until spring

3 kinds of continued fever viz. Synocha Typhus & Synochus Synocha char by a phlogistic state of system throughout the whole course of disease When local inflammation commences with the fever then it is called phlebotomy or phlegmosis Typhus exactly the reverse small & peg pulse depression

Continued Fever, February 3rd 1837

Synochus constitutes the predominant fever of our winters
February 6th 1837

Synocha

It often commences very obscurely & ^{is} slowly developed
At other times it comes on abruptly

If the fever lasts only 24 h it is called Ephemeral
Cause There does not appear to be any predisposing cause

Atmospherical changes sudden exposure to cold
debauchery sudden effects upon the mind

It is the most simple & least dangerous form of
continued fever

In the treat we have merely to keep in mind the
nature of the fever & the ease with which the fever
is broken up

Diet & regimen, diaphoretics, Purgatives you
have the effect of curing this form of fever

Typhus

It is an infectious disease depending upon
exhalations arising from ^{from} body Diff: from contagious
in this that contagious diseases are produced from
persons labouring under the same disease & at
the same time being incapable of being produced in
any other way

Feb 7th The symptoms are held together loosely
in Synocha & is to be treated with mild remedies of
origin of typhus It freq emanates from persons labouring
under typhus It also occurs where ^{heavily} persons are crowded
together Can it arise from any other cause than
animal exhalations arising from morbid disease?
More animal putrefaction will not produce ^{the} disease

Typhus.

February 7th 1834

strict ventilation night & day merely considering the comfort of the patient. Allow no curtain about the bed & the patient should be placed in a spacious apartment. ~~Persons~~ should avoid inhaling the patient's breath. avoid currents of air from patient's person & body cloths should be freq. changed & when removed immediately plunged into cold water.

Disinfecting agents Vinegar Nitric Acid
Chlorine gas is more efficacious than nitric acid but not so disagreeable.

Take Nitrate of potash & pour upon it sulphuric acid when Nitric acid fumes will be liberated.

I think it is to be prepared in sick chambers. High temperature is unfavorable to the production of typhus from infection. It occurs most freq. in cold weather.

Symptoms

They are very variable. It is from this circumstance that authors have attempted to establish diff varieties.

I shall not adopt the modification of Armstrong.

Stage of depression Stage of Excitement
& stage of collapse

1st Stage of depression. Comes in grad manner. lasts 4 or 5 days. ~~consciousness~~ ~~disagreed~~ ~~mind torpid~~ ~~countenance~~ ~~dull pale~~ ~~alternating~~ ~~sensation of cold~~ ~~in~~ ~~temperature~~ ~~very irreg~~ ~~tongue~~ ~~white~~ ~~relaxed~~ ~~with marks of teeth~~ ~~appetite~~ ~~lost~~ ~~very little~~ ~~thirst~~ ~~pain in loins~~ ~~oppression~~ ~~in precordia~~ ~~pulse small & feeble~~

2nd Stage. Grad develops itself. pulse is grad developed. appetite totally lost. rare cases there a craving for food. vomiting & thirst most day & parched. exacerbaton comes on about 6 or 7 P.M. delirium comes on sometimes constant burning heat.

towards morning the patient is rather more comfortable
 & patient takes a short sleep. The signs
 continue ~~for~~ this way for about a week.

3rd stage comes from 8th to 11th day
 The fever has ~~ceased~~ Great fetor in all the
 exhalations great inaction with ulceration ^{there is prostration} in parts where
 pulse is diminished in force & increased in frequency
 patient is very weak scarcely able to move or speak
 tongue is dry & stiff disposition to slide down
 in bed Sordes is formed Sensorial functions
 are failing There is a low muttering delirium & various
 visual allusions as pushing bed clothes
 are the evidence of great prostration is twitching
 of muscles & when affecting face subsultus tenetum
 diff deglutition involuntary discharges from bowels Coffee ground
 vomiting &c

præticæ merely look as if perished upon skin
 & resemble maling

Feb. 8th 1837

This stage cannot long continue
 The power of voluntary motion is lost hypochratic countenance
 eyes turned laterally & superiorly respiration is hurried
 & panting & after a while the respiration stops &
 spasmatics become relaxed

There is no disease in which there is so close an approx
 between life & death as in this

Cerebral inflammation is the most preg ~~inflammation~~
 that occurs in typhus

The tips of the lungs not so liable to be involved
 The texture of the digestive organs are very liable to be
 involved

Therelated inflame do not uniformly occur in typhus
 typhus attended with angustia
 Pathological Anatomy

Typhus

February 8th 1837Pathological Anatomy -
in glands of the

ulceration of last part of ileum

They occur in course of fever

They sometimes penetrate

the peritoneum
February 9th 1837

Synochus

Can continue perch of all parts of this Country
most common during cold & variable weather though it
occasionally occurs at all seasonsAmong our Physicians there is no distinction
made between this & typhus

They differ in origin

Synochus arises from
cold variable damp &c. while typhus arises
from infection

Treat. of Synochus & Typhus

Is it in the power of art to cut short Typhus

It cannot be done abruptly

It is possible at once to stop the course of Synochus
We can mitigate the symp of typhus - Predominant
feature is prostrated the systemIn 1st stage we are to relieve the oppressionIn 2nd - to relieve excitementIn 3^d stage to prevent the system from sinking1st Stage of TyphusTo bring about reaction is our 1st objectPlace patient in spacious apartment 1st by

Rarely advisable using powerful stimuli Carbonate of ammonia &c
 to practice bloodletting after 2nd or 3^d day of
 stage of fever
 But is it a proper remedy? In many cases it is not
 when we can possibly avoid it
 Best means is an emetic in my opinion
 I consider Z. Emetic the very best remedy
 occasionally I prefer it the best

Typhus

February 9th 1837

When you wish to produce emetic, cathartic effects
 2 or 3 grs. Z. E. with 15 or 20 grs Calomel
 It is desirable to produce copious emesis. Reaction
 gen. shortly fol. The disease is frequent short
 here

2nd Stage or Stage of Excitement

Whether synochus or typhus
 Blood-letting - can consider a prep rem It should
 not be liberally employed - but merely to relieve
 the stage of excitement In latter period of this
 stage it is a doubtful remedy

Must adapt local rem according to circumstances
 If inflammation of brain exists - cold applications

to head cupping counter irritation &
 Purgatives in 2nd stage 4 or 5 stools daily = Armstrong
 I consider it improper carried to that extent
 Calomel alone or combined I prefer a free
 dose of cal alone fol by Castor oil & prep. genna
 in equal proportions

Cold Effusions I have seen great benefit derived
 from it It should be used at night during
 exacerbation When temperature not above
 natural standard When there is chills from
 striking patient it should not be used

February 10th 1837

3rd Stage or Stage of collapse

Carbonate of Ammonia nourishing food diffusible
 stimuli. sinapisms to epigastrium

As patients convalesce, should be exceedingly cautious in regard
 to diet Liqueur mucronatus morphine about
 the strength of Laudanum = to be used in cases
 of watchfulness in typhus

Proximate Cause of Idiopathic

Fever

February 10th 1837

Def. Example Gastric indigestion by ~~pep~~ morbid sensibility
 thirst & ~~etc~~ Inflamm is in this case the proximate cause
 Do we know the proximate cause of Idiopathic fever?
 Is the proximate cause always the same?
 Hypothesis of Clutterbuck ~~Brady~~ & Fordyce
 Clutterbuck attributes the proximate cause of
 fever to inflammation of the brain

February 21st 1837

Exanthematous Diseases Feb 21st

Rhazes 1st described small pox. It existed in the 10th
 Century Patient affected with small pox ~~will~~
 is capable of producing the contagion from the time
 that the eruption 1st makes its appearance until the ~~last~~
 scab falls off
 Exanthemes diff from idiopathic fevers in certain particulars

Notes taken from Dr. Green's Lectures on Chemistry

Nov 5th

Matter is the sensible substance which composes all bodies

When any body is separated into its elementary principles it is said to be analyzed

Matter exists in 4 forms Solids, Fluids, elastic fluids or gases exist free in the atmosphere. All known elastic fluids are transparent.
4th Ethereal or unponderable substances. Heat light electricity & magnetism
Term element is synon. with undecomposed body

Synthesis consists in forming a compound by joining its elements to unite

Gravitation

Crystallization

1 The Parallelopiped the simplest of the solids whose faces are six number & parallel 2 p. 2

How is prussic acid made & how is it detected in the stomach of a dead person? 153

Made by heating cyanuret of mercury in a glass retort with $\frac{1}{2}$ its weight of concentrated muriatic acid

Free hydrocyanic acid is easily recognized by its odour. Chemically it may be detected by agitating the fluid supposed to contain it with the peroxide of mercury in fine powder. Water & the cyanuret of mercury are generated & on evaporating the solution slowly the latter is obtained in the form of crystals

Prussic acid made by passing sulphuretted hydrogen through a solution of the cyanuret of mercury

Arsenic

Jan 10th 1837

Does arsenic preserve the body? It does act as preservative

Animal has been preserved for 2 or 3 yrs

W. Christ Test

Take out stomach & tie up the 2 ends & take it to your laboratory or some other place where no other person can have access

Turn out the content of stomach into a clean dish & then examine the coats of stomach & if you discover a white particle place it on a piece of red hot charcoal. Thereupon Carbonic acid will be formed & a disagreeable odour will be

144
Arsenic

Jan 12th 1837

If you have the aliacous odour then you may be sure that arsenic exists. If none of these white particles are discovered you proceed as fol. Dry the matter & divide it into parts. Rub ^{small portions of} 1 part up with charcoal & then throw it upon coal. With clean Florence flask put the portion of matter & pour upon it distilled water then boil again & again. Throw the matter upon filter & filter it when hot. Solution, - fluid tests. Ammoniated sulphate of Copper dissolved to get exact proportion.

It produces a wheely green. It corroborates the other test. The 2 evidences renders it certain. This as a test taken alone is not sufficient proof. Ammoniated Nitrate of Silver produces yellow colour.

Jan. 13th All the arsenites are soluble in fluids of stomach. Lime water is not a good test. I shall not say any thing about it. {which is yellow}

Ammoniated Nitrate of Silver. Arsenite of silver is precipitated =

All the above tests together prove that arsenic exists. Sulphuret of Antimony pour upon it. Muriatic acid to form. Fluid supposed to contain arsenic must have some acid put into it = muriatic acid & then pass through it sulphuretted hydrogen when orpiment will be formed = sulphuret of Arsenic & yellow.

Another Method. began on this day take the matter dry it & make a solution with distilled water. Pass through it sulphuretted hydrogen. 2nd apply heat & drive out sulphuretted hydrogen & sulphuret of Arsenic will be deposited. filter it & put it in a glass tube larger if quant be great smaller if the quant be small. Put upon it black flux made by dephlograting. Take Co Potash or soda drive out the water of crystallization & mix it with charcoal when you have a black flux. Apply heat. And the arsenic will be driven in form of white oxide of arsenic. Above it being

Isenic Jan 13th 1837

Antidotes Sulphuret of Potash no antidote
 Magnesia It cannot act as one
 Charcoal is quite as inert
 Hydrated oxide of iron = Sulphate of iron into nitric
 acid digest them together & throw in ammonia
 very little dependance can be placed upon it
 Best way is to give an emetic = Sulphur Acid Take diluent
 drinks stomach tube injurious
 Milk is the best antidote by coagulating in stomach
 envelopes the particles Saccharine matter is
 also a very good remedy
 January 16th 1837

Stannum

It is found in the state of Peroxide.
 It does not oxidate but tarnishes when exposed to air
 Peroxide made by pouring dilute N Acid on tin foil
 Tin taken internally is dangerous because it is apt to be converted into oxide
 Peroxide of tin fused with glass forms enamel
 Test Sulphuret of hydrogen gives in a yellow precipitate

Jan 23rd

Potassium

Lower fifth of earth composed of magnesia lime silica alumina
 Protoxide of Antimony it is formed in process for forming sulphuric
 hydrogen It is the only active prep. of antimony
 1 atom of each
 Protoxide formed by pouring nitric acid on Antimony
 Sulphuret is the ore from which we derive the metal
 Hermin Mineral

Phosphore of Antimony & P of Lime - James powder
 made by 2 parts horn shavings & 1 part sulphuret of
 antimony & apply heat Protoxide is the efficient
 remedy

Ammonia is the test for copper It produces a blue colour
 Copper in almost all its salts may be detected by dipping a piece
 of polished steel.

Coin is 23 carats fine

Red precipitate detected in fluids of stomach by adding proto.

side of tin when mercury is revived

The sulphuret of mercury = Vermillion Its presence may be detected by combining it with potash the mercury is revived

To detect Corrosive sublimate

1st make proto chloride of tin

Oxalic Acid after evaporation for Sulphur magnesia

It proves suddenly fatal in dose of an oz or more
Antidote Bicarbonate of soda lime magnesia

150

Notes taken from Dr. Pattison's

1836

Nov 14th

Lectures on Anatomy

Foramina of Temporal Bone

3 classes - 1st class those which perforate the bone and belong to the organ of hearing

2nd class those for the transmission of Nerves

3rd class for the transmission of blood vessels

1st Class Meatus auditorius externus, iter ad palatum & auram, Aqueductus vestibuli, Ag. cochlea

2nd Class Foramen 2 upon posterior petrous surface

1st Foramina Meatus auditorius internus which transmits Portio dura & portio molis And in this is the aqueduct of Fallopius

2nd Anter petrous foramen or vidian foramen & the stylo mastoid foramen by which portio dura passes out

3rd Class Foramen of Spinalis only other foramen is just behind mastoid process & is sometimes wanting: it transmits small vein & opening is common to this & to occipital foramen

Lacerum posterius Transmits 8 jugular & pair nerves

Ethmoid bone It is an oblong square fig. It is wedged between the 2 orbital processes

Optic Foramen opticum At lacrum orbitale Foramen lacerum medium sit between sphenoid & temporal bones

3rd Foramen division of 5th & 6th nerves pass out of foramen Anter

5th Foramen vasa optica venae

1st Foramen Foramen rotundum immediately behind foramen Lacrum

2nd Foramen Foramen ovale sit little farther back

3rd Foramen Foramen spinosum 2nd branch of 5th pair goes out of foramen Rotundum

4th Foramen Foramen transversum all over of entrance of A. Meningia Media

5th Foramen Foramen ovale

6th Foramen Foramen spinosum

7th Foramen Foramen transversum

8th Foramen Foramen spinosum

9th Foramen Foramen transversum

10th Foramen Foramen spinosum

11th Foramen Foramen transversum

12th Foramen Foramen spinosum

13th Foramen Foramen transversum

14th Foramen Foramen spinosum

15th Foramen Foramen transversum

16th Foramen Foramen spinosum

17th Foramen Foramen transversum

18th Foramen Foramen spinosum

19th Foramen Foramen transversum

20th Foramen Foramen spinosum

21th Foramen Foramen transversum

22th Foramen Foramen spinosum

23th Foramen Foramen transversum

24th Foramen Foramen spinosum

25th Foramen Foramen transversum

26th Foramen Foramen spinosum

27th Foramen Foramen transversum

28th Foramen Foramen spinosum

29th Foramen Foramen transversum

30th Foramen Foramen spinosum

31th Foramen Foramen transversum

32th Foramen Foramen spinosum

33th Foramen Foramen transversum

34th Foramen Foramen spinosum

35th Foramen Foramen transversum

36th Foramen Foramen spinosum

37th Foramen Foramen transversum

38th Foramen Foramen spinosum

39th Foramen Foramen transversum

40th Foramen Foramen spinosum

41th Foramen Foramen transversum

42th Foramen Foramen spinosum

43th Foramen Foramen transversum

44th Foramen Foramen spinosum

45th Foramen Foramen transversum

46th Foramen Foramen spinosum

47th Foramen Foramen transversum

48th Foramen Foramen spinosum

49th Foramen Foramen transversum

50th Foramen Foramen spinosum

Processes of Sphenoid = Great wings or temporal processes

Spinous process, Styloid process Transverse spinous process, Trigonal process, Zygomatic process, Olfactory process, Triangular process

Anatomy Nov. 27th 1836

Foramina of temporal bone Meatus Auditorius externus
 Meatus Auditorius internus by which the auditory nerves
 have access to ear The Meatus Fallopius through which
 the posterior dura passes Median foramen sit on anterior
 surface of petrous portion of temporal bone It receives a
 small twig reflected from 5th pair of nerves
 Stylo mastoid hole by which posterior dura passes out
 After a plate ad arium or Eustachian tube by which
 air enters into ear

Carotid foramen enters skull near point of petrous portion of
 temporal bone Sympathetic nerve is also transmitted
 through this great canal

Foramen lacerum posterius formed by temporal & occipital bones
 It transmits Jugular vein & 8 pair of nerves
 Mastoid foramen Ducts of Cerumen

Processes Zygomatic, Styloid, Vaginal Mastoid Auditory processes
 December 7th

Larynx

Connection between Thyroid cartilage & Hyoides

Thyro hyoid membrane connects thyroid & through whole extent to
 hyoides Also connects by lig. extending from posterior
 horn of Cartilage of Hyoides

Cartilage of Hyoid & horns 2 posterior superior & 2 poster. inferior horns
 Cricoid Cartilage is a ring broad behind & narrow before
 Cricoid Thyroid membrane is the space through which opening for Larynx is
 Aryepiglottoid Cartilages small triangular shaped cartilages
 attached to posterior part of base of Cricoid

Epiglottis attached to base of hyoides Pituitary opening into
 wind pipe Artery Epiglottidis

Dec 14th Muscles of Neck

Superficial muscles of side of neck

- 1 Platysma Myoides It supports vessels of neck & adjusts them in circle
- 2 Stern. Cleido. Mastoideus 2 heads become connected about middle of neck
 It is 1 of the land marks for surgical anatomy of neck It is enclosed in a
 sheath formed by layers of deep fascia of neck Its position can always be
 seen in living subject

2nd Division Muscles sit. between lower border
of os hyoides & Sternum

Sterno Hyoides origin comes out from chest

lay. like a rib on fore part of Trachea These 2 unite in middle
by cellular tissue completely concealing trachea. On removing this
muscle I bring into view the

Sterno Hyoides it arises from inner part of sternum & from
cartilages of 1st & 2nd ribs. It lies rather to outside of above muscle.

It does not lie in contact with muscle of opposite side

Omohyoides has 2 bellies arises from notches of scapula
tapers in course Under I.C. mastoides, it is tendinous

It goes up along side of stern. Hyoid & is inserted into junction
of body & horn of os hyoides

Thyro Hyoides It lies under before we can open thyro Hyoid
mem. we must separate these 2 muscles

Cricoid Hyoid M. does not divide these muscles in joining mem.

They leave & open space triangular

Thyroid Gland cov. by ^{sternohyoid} Thyroid, Laterally by ^{thyroides} Omohyoides

3rd Division Muscles sit between chin & os hyoides

Digastricus. arises from mastoid process. It
goes downwards & forwards to body of os hyoides
passes through band of Cooper & then proceeds to chin

1st of Hyoid muscles Stylo Hyoides inserted into junction
It lies above Digastricus

Stylo Glossus

Stylo. Pharyngeus opens & raises the pharynx

Neck divided into upper middle & lower region

Middle reg. bounded by 3 muscles Laterally by outer border

of I.C. mastoides. Above by lower border of Digastricus

Anterior. ~~Super~~ by super border of Omohyoides

All lying below this is the lower region All above the super

This triangular middle space the ~~parts~~ are imperfectly
muscles. A line from deapression of I.C.M. directly upwards

describes the course of Cerebral artery Connections

It is enclosed in sheath with preceding artery nerve & enters Jugular
artery lies next to trachea nerve in middle vein to outside

& helps directly through middle of middle region of neck
It is hidden upon longus colli at lower part & at upper

Muscles of Neck Dec 14th 1836

On dividing skin facia & platys myoides the vein pt comes into view
formed by junction of jugular, temporal

Exter Carotid passes behind Digestricus & Stylo Hyoides & before
Glossoglossus & Stylo Pharyngeus

Hyoid muscles coincide in pulling tongue back & pulling throat upwards

Dec 15th Muscles sit between Hyoides & Jaw bone

1st. 4 in number

1. Mylo Hyoides To obtain a full view must remove anter belly
of Digestricus Arises from mylo Hyoid ridge inserted
into upper margin of Hyoid bone

2nd Genio Hy. immediately under mylo Hy & therefore from
3rd layer It from chin & inserted into body of Hyoid bone
Arises under central origin of Mylo Hyoid

On removing these there is another layer in the 4th layer formed by
3rd Genio Hyo glossus It arises by pointed origin from chin &
inserted into

Another in Hyo glossus It arises from body of appendix & inserted into
inserted into side of tongue Tongue receives Hypoglossal nerve
for muscular motion Gustatory sense of taste

glossopharyngeal respiratory nerve
Lingualis Muscle between fibres of genio

Action is to contract the substance of tongue

Dec 15 Hyoid bone connects directly to trachea by Thyro Hyoid & omohyoid muscles
Longus Cilli muscle lies anteriorly on sides of bodies of vertebrae
Carotid artery lies pillared upon it
Rectus Anterior major occupies same sit, but higher than longus
minor

Muscles along side of neck

Scalenus Anticus comes up from under corner of clavicle
Behind this there is a large bundle of fleshy fibres which is
Scalenus Posticus Arises from 1st & 2 ribs

Muscles arising from bones of trunk & inserted into bones of superior extremity

Pectoralis Major

Ligaments of knee joint. Strength of joint is confined to the ligaments
1 Internal & 2 external lateral ligaments. Strong bundle behind
arise from back part of external condyle of femur = Posterior ligament
inserted into head of fibula = Ligamentum posterum Winslow

Ligamentum Patellae strengthens articulation upon its anterior face
Crucial ligaments. 1 arises from external condyle and is inserted
into pit before the eminence upon tibia. 2 arises from internal
condyle & inserted behind

Ligament alaria major & minor are seen upon opening the joint
upon each side of patella. Ligamentum mucosum
flexion joint upon the back of the posterior ligament

Relations of Sciatic nerve to rotator muscles of thigh
on removing Gluteus Maximus the Sciatic nerve is exposed
It comes out below Piriformis & crosses the gemelli superior obturator
internus Gemelli Superior & quadratus femoris

It runs down between flexor muscles of leg with semitendinosus on
one side & semimembranosus on the other

On amputating thigh at hip joint you divide the rotators of thigh
On removing quadratus femoris you bring to view the obturator
externus which is a powerful rotator of thigh outwards

Jan 16th. 1887

Arteries — 1st Carotid

Middle cervical region of Neck is a triangular space bounded
laterally by ^{upper} border of Sternocleidomastoides Inferiorly
by upper border of Omohyoides Superiorly by lower
border of digastric muscles

Artery runs through the middle of this triangle. In cutting
down upon it you merely divide the ~~superficial~~ superficial
facia Platysma Myoides & deep fascia of neck

Surgical Relations. Between artery and jugular vein
lies by side of carotid & in common sheath with it
between artery & vein you have a large nerve & intimately connected
Pneumogastric nerve is the great nerve of respiration & digestion
Great sympathetic lies behind the artery & only separated from it
only by the arched sheath which encloses the jugular Pneumogastric
nerve & carotid artery

Jan 6th 1837 Common Carotid Artery

Divides at upper border of thyroid cartilage
 Sometimes the carotid does not divide until it arrives at angle of jaw
 In some rare instances the 2 arteries arise from it

Jan 11th 1837 Skin superficial fascia composed of 2 laminae
 between which Platysma My. is formed Deep seated
 fascia much stronger The laminae form a sheath for ~~arteries~~
 glands mastoid One lamina goes upwards over external surface
 of parotid gland & goes over face The other lam goes behind
 parotid is attached to styloid process &

It is attached to posterior transverse process of 6th vertebra
 One lamina goes before & the other behind the ~~artery~~
 sheath surrounding the artery when open we discover the Pneumogastric
 nerve lying on outside of behind the artery Operation of
 tying carotid it is an easy operation

Glandula concatenata very freq the seat of disease they
 arise from angle of jaw to lower part of neck & closely connected
 with carotid

Relations of carotid in lower region of neck It has precisely relations
 same regard to vein & nerve

Stems C. Artery Thyroid & Hyoid cover it

Diff between arteries of 2 sides Right is the shorter
 Artery of left side is sep from trachea by gullet
 In cutting down upon oesophagus run very little risk of wounding
 the artery there is a danger of wounding nerve which lies
 upon anterior surface of gullet - recurrent Laryngeal nerve

Branches of carotid 1st into external & internal
 External 2nd & 3rd 1st before forwards
 2nd backwards 3rd

1st Clap of branches going forwards

Superior Thyroid arises immediately after division of carotid
 It gives off Super laryngeal artery which perforates the thyroid
 gland main Super laryngeal nerve lies immediately behind
 the artery 2nd course down & forward goes under omohyoid
 muscle to thyroid gland

3 nerves go to tongue *Glossopharyngeal*

Jan 21st 1831 *Glossopharyngeal* ^{connected with lower border} ~~is upon lower~~ of *Stylopharyngeus*
muscle
Internal carotid goes up neck ^{passes above} ~~sectes~~ Anterior
4 papers behind *Stylo Glossus* & *Stylopharyngeus* &
she interposes between the 2 carotids
Digastricus & *Stylo Hyoides* cross external carotid
externally. It

Jan 26th Relations & Connections of The Subclavian
& Axillary Arteries

connections of
Arteria Innominate crossed by Vena Transversa
cov. by S.C. M. & H. & Thyroid must be detached from
sternal origin to tie the artery
There is danger in tying external Innominate
Subclavian Artery. R. V. M.
1st cardiac portion S.C. M. St. V. Sterna Hy.
cover it. It is cov. & concealed by confluent
of Subclavian & jugular. *Def. Pneumoz* N lies upon
it anter. surface. Encircled by Recurrent Laryngeal
which goes of from *Pneumogestric* N crossed by lower branches
by ganglionic nerves. Likewise crossed by *phrenic* nerve
much more diff to tie this artery than innominate.
It would be the most diff operation in surgery
Left Subclavian Cardiac portion
diff between 2. Right comes of secondary from
artery

sit much deeper on left side than right
Artery of left is cov. by carotid & by vena transversae
2 gastric & *phrenic* Recurrent Laryngeus Thracic
duct empties into junction of Left Subclavian
& jugular
No man in his senses would think of performing the
operation of tying ~~the~~ artery

Connections &c of the artery in 2nd part of
Subcl. vein lies behind the clavicle sep from artery by scalenus
anterior muscle

Last Portion of the artery

Axillary plexus lies connects with artery & is the only important
connection. There is no reason why the artery

may not be tied in this situation except where there is aneurism
in the axilla where the clavicle is carried high up

It is much easier than tying the axillary below clavicle

Axillary & Relations at upper part

Ar vein below nerves above

diff to tie the artery upon separating the 2 portions of
Pectoralis major you have the Artery before it
Coraco clavicular lig also interferes

Lower Portion

It has vein lying below
& also 2 nerves &c

Relations & connections of Anter tibial artery after perforating
the inter. membrane it sends of recurrent branch

At anter part of leg it lies deep lower down more superficial

At upper part of leg it lies between Anterior tibial muscle & exten
sor dig comm lower down it lies between extensor

of great toe and anter tibial muscle

At ankle it furnishes 2 alveolar arteries then passes
under the annular lig. to foot gives of tarsal artery

It lies superficial just towards root of metatarsal bone of
great toe but before it passes it sends of the

Posterior tibial artery. Reached by sep attachment of head of Soleus
from inner side. It lies between Posterior tibial &

femor dig communis. It is intimately connected with nerve

It passes in hollow between malleolus internus & os calcis with
connected with nerve. It divides into ep & inter plantar arteries

162

165

Notes taken from Dr George McClellan's Lectures on Surgery

1836

Nov. 15th Shortest period in which I ever knew ~~an~~ inflammation overwhelm to take place after an injury was 12 hr. G. McClellan says shock In other cases it does not take place for several days

Insidious Shock

Most dangerous period is the period supervening between the injury & the inflammation in severe injuries Surgeons are very apt to be deceived in this case the patient answers questions & every function appears to be natural The patient lives but a few hours

Case of Dr. McClellan

I have seen many such cases - G. McClellan Ankle joint torn open bones protruded & perhaps the blood vessels &c Symp. The functions all appeared natural for 2 or 3 days when I was called. She had a peculiar expression of countenance with a degree of surprise She died on 3rd day It is like the injury to a ship when the anchor is cut & the ship glides on peacefully to a wharf while all are crying peace until &c

Nov. 21st

Morbid Inflammation

Debilitated persons liable to Morbid inflammation Local inflammation when attended with debility should be treated by gen local & gen stimulants

Erysipelatous Inflammation When you come beside the patient you examine into the state of general sys. & particularly the state of pulse

Bleed as frequent as the case may require I bled a drunken woman, who had been kicked down stairs by her husband and received an injury of scalp from teeth of comb which was followed by erysipelatous inflammation, 10 times in the course of 24 hours & she perfectly recovered

Nov. 22nd

She must have lost 60 or 70 of blood * 170 Irritable pulse known by the previous history of case, by the patient becoming worse when raised from pillow by great throbbing of the carotids at the same time that they may be compressed by moderate pressure Face is flushed there is agitation & delirium

170

Surgery Nov. 22nd 1836

* Continued from 169. In patient with phthisis pulmonalis even in last stage if pleurisy or peripneumonia supervene you must bleed & use other antiphlogistic remedies.

Nov 23rd Morbid Inflammations & etc

Erysipelas

Treatment - Bleed, purge, use Antimonials demulcent drinks & apply mucilaginous and other soothing applications to the parts. Sugar of lead water &c

When the inflammatory symptoms run high you must bleed largely & continue to employ it until the pulse is reduced & the inflam symptoms moderated

When from long continuance of disease or when it occurs in debilitated subjects it may require moderate stimulants. Whey is the very best that can be used. It should be made in fol. way. Take 50 pt. new milk boil it & throw in while boiling from $\frac{1}{2}$ to 1 gill of some of white wines such as ~~Port~~ Madeira &c & afterwards strain it & sweeten it with loaf sugar.

If this is not sufficiently nourishing you may add to the whey & equal quant of Sago or Tapioca boiled so as to make a thick mucilage & sweeten it with loaf sugar.

Blisters turpentine corrosive washes nor any other irritating substance should not be applied.

Blisters may be advantageously used when the inflam is spreading by applying 4 inch or two above the inflam surface.

When matter forms in sub-cutaneous cellular tissue you should make incisions so as to coacenate the matter.

Nov 24th Phlegmonous Erysipelas requires active antiphlogistic treatment. Incisions into the part & poultices &c

Punctured wounds. When person receives a wound from dissecting dead bodies the inflam is apt to extend up the arm so that it becomes enormously swelled. It will be followed by high symptomatic fever. It should be treated by copious bleeding purging & antimonials calomel & opium.

Local treat

171

Morbid Inflammations Nov. 24th 1836

Specific poisons which are produced from bite of adder, wasp & other poisonous animals act upon the system differently from punctured wounds. Instead of its being fol by great local inflammation & symptomatic fever. The effects of the poison is upon the whole generally attended with great tendency to sinking in system & instead of the system requiring depleting remedies it must be supported by stimulating remedies such as Aqua ammoniac spirit of ammonia carbonate of ammonia.

Local treatment. We should endeavour to prevent absorption of poisonous matter by exposure above the part scarification & suction.

Nov. 25th You should always graduate your remedies to the state of system. Phlegmonous & Anthrax. They mostly originate from some internal cause. They both have the same tendencies. Anthrax very seldom requires stimulation other only when patient is phlegmic. The system is more frequently depressed & requires depletion. It is sometimes necessary to bleed 4 or 5 times.

Local treatment. Leeches, cooling & moist applications. & free incisions into part. I would not mix with acids may be sometimes used as escharotic with advantage to sloughing part.

Mortification — It is in all cases attended with prostration of vital powers. It is generally owing to high inflammatory action.

Nov. 28th In every case of injury attended with mortification where I have examined I have found matter beneath.

Treatment It will require antiphlogistic treat until the inflammatory symp. have been reduced to a proper grade of action & where matter is confined free incisions must be made.

Local Applications Linen rag wet with mucilage of elm bark should be applied to the mortified & inflamed parts.

To destroy the feta may apply Chlorate of soda charcoal or diluted pyrolygneous acid.

If constitution flags give stimuli such as wine whey with tapioca sago arrow root mucilage or if it requires more permanent stimulus give brandy & water. Essence of beef which is made by taking lean part of an ox or some animal put a pound of it in a bottle

& boil it for $\frac{1}{2}$ or three hours &c

Mortification Nov. 29th 1836

Basilicon ointment may sometimes prove useful by stimulating the surrounding parts. A few drops of Sp. Turpen. may be added.

Mortification occurring in old persons best treated by giv. solid opium & combining it with nourishing diet. Sometimes necessary to combine the above treat. with bloodletting. Apply mild & irritating applications to parts = mucilage of sugar of lead water.

Gangrene from pressure change position of patient employ mild applications lead water, mucilage, poultice after inflammation is subdued apply litharge plaster.

Abscesses

Nov. 30th Abscesses which do not point We frequently meet in such cases where punctured wounds have been inflicted. Great constitutional irritation is produced by confinement of matter beneath tendinous structures. Surgeon should never wait for the pointing of the matter but evacuate the matter.

Mode of opening the parts Make a free incision by a scalpel or if abscess be very large make more. Select the points which your judgment may indicate. Afterwards place soft elastic bandages around limb so as to press the matter out & apply poultices upon limb. Should make cuts in bandages over incision.

By this means I have seen them healed in course of 2 or 3 days. Constitutional treat Use a light & irritating diet and use other antiphlogistic treat.

When abscess occur in peritoneum it is gen. attended with serofulous symptoms.

Injuries or bruises attended with extravasations of blood may in healthy constitutions be absorbed by bleeding purging. Low diet antimonials discontinue when shivering of summer.

Cold Abscesses They may be formed without inflammation occurring. They occur in persons of debilitated constitutions & rarely occurring in healthy looking individuals.

any many
 Abscess Nov. December 1st 1836

They should be opened by a calvarial incision in edge of
 of tumor & a probe introduced so as to allow the matter to
 flow out. But you should not press it. Afterwards
 a dip of lint & over that a strip of adhesive plaster
 & keep the patient at rest. If the inflammation has a tendency
 to run too high apply lotion of lead water vinegar & water
 muric acid of ammoniac &c.

Constitutional Treatment Patient should avoid all
 indigestible articles of diet. Should use stale wheaten bread
 sage arrow root rice barley until tone of stomach is restored
 when patient may use animal food &c.

To correct the secretions, of the intestinal canal especially must give
 alteratives. Magnesia. when acidity prevails.

December 2nd After the abscess has been opened the matter which
 will remain may be absorbed by the above Constitutional treat

Burns Local Effects

1st kind where the effect on skin is transient

2nd — where cuticle is raised in form of blister

When burn extends through the rete mucosum there will be
 troublesome ulcers. When the burn extends still deeper

we invert the ratio of dangers — they do not suffer so much

as in former variety. The most dangerous period ^{is when} discharge

of pus takes place

Treat should ^{use} soothing remedies. Where blisters have not been
 formed should endeavour to prevent them & when they
 form you should be careful so as not to remove it but
 make small punctures.

Local Remedies Laying 1 or 2 layers of cotton over any common
 burn is the best application. It prevents violent of
 temperature.

2nd raised or benighted burns should evacuate serum by small punctures.
 Cotton is still the best rem. when cuticle remains.

If part become constricted pour whiskey brandy & water
 or alcohol on the cotton. Any kind of soft or undisturbed
 applications may however be used.

174
Burns. December 2nd 1835

when cuticle has been removed should prevent irritation of air &c. Where there is a large surface exposed cotton moistened by ^{dilute} alcohol is the best application at same time, should use constitutional treatment. Aquafortis & Ammonia

Dec. 5th is also a valuable application. Slightly diluted Strammennund ointment is a very good application in an abraded burn when it is the most very irritable Elder ointment &c answers very well.

Linseed boiled down with sublimed oxide of lead to the consistence of a plaster is as good an application

as can be employed when abraded surface is not to irritate.

Shoe-burns

Efficient Poultices are the best applications that can be used. Constitutional disturbances of burns. When they are large we have an overwhelming shock. The patient becomes cold insensible & dies.

Another kind of syn- sometimes occur. The patient will cry & moan & then they will become cold & prostrate.

When the pulse becomes full skin hot & the reaction is high the constitutional will not bear the effects of depletion as well as in other inflammations. But in such cases

(inflamed fever) depletion is necessary to moderate symptoms. Cool diluting drinks. Laxative injection. Saline laxatives.

Sponging the undrained surface as in other fevers.

When I get the skin perspiring then I have hopes.

Sometimes the system will not rally or if the system does rally

in some degree it relapses again. In such cases the Lentish ointment is very serviceable.

Dec. 6th

Effects of Low Temperature.

There is a loss of blood in external parts while internal parts are congested.

Letter

You must tend to the system. Avoid damages. Pasteries heavy & indigestible articles of diet.

very often indeed you will fail unless you confine your patients to the lightest articles of diet. Use of diluent drinks. Diaphoretic. Gum arabic.

175 30 33

Surgery December 8

perfect cleanliness alteratives tongue is foul beneath
 is foul $\frac{1}{2}$ gr cal or 3 gr blue pill every night together
 with antimonials with diaphoretic drinks Warm bath
 If there is a specific in Herpes Arsenic is that rem in
 dose of from 5 to 8 gr 3t, d.
 Local Rem 1 to 3 gr in oz water or nitrate of silver
 in solution

They are the best local rem that can be employed

Note the tongue on eating cancer
 sometimes begins in small pimples It occurs on back of hands must
 frequently on face nose back &c
 They are best removed by caustic Arsenic applied locally is
 very good best oint 3

10 gr arsenic made stiff by
 litharge plaster & spread it
 on hind skin

Scrophula - Dec 9th 1836

It is a disease of whole system It affects bones
 muscles tendons skin as well as glandular system
 Most inveterate form is where it is hereditary
 The lungs are deperant in organization The chest is contracted
 There is a dulness of percussion under scapula
 Tubercles are from size of millet seed to that of hazel nut
 Scrophula also develops itself in glands of neck, mesentery
 when scrophulous sys occur externally tubercles seldom occur
 in lungs

Acquired form of Scrophula

Insufficient food & clothing & where they are brought up
 in crowded rooms Also in children of the most luxurious
 & wealthy Treat of Plered Scroph

cannot perfectly cure we can not save the part to be destroyed
 Regulate diet Abundant exercise Sufficient clothing

176
Scraps December 9. 1868

Dec. 9th

Sec. 9th Regulate the state of the bowels
Patient must use exercise in open air even in winter
Nourishing food avoiding stim when fever exists
Use exercise on horse back for $\frac{1}{2}$ h at a time

The treatment of a young Lady who was suffering under the most malignant form of Scrofula Under the treatment of J. The first day that the young Lady was placed under his treatment he had her to ride on horseback though she was so much exhausted that she was under the necessity of being held on the horse while the horse was lead about forty length of year hour she was then taken from off the animal quite much exhausted the next day the young Lady went through the same process of exercise ~~only~~ with an interval of time 5 or 10 minutes and so on taking this exercise for some length of time she at length recovered in strength sufficient to ride the distance of a mile or more with any assistance still increasing in strength at last she recovered so far as to be able to ride the distance of 5 or 10 or 15 miles she at last set out on a journey of some distance across the mountains to visit her relations which mountains ^{she} crossed three times though the course of the winter and ~~now~~ I am happy to say that she is now one of the most amiable & healthy woman in this City with the only exception of her shoulders are somewhat contracted and truly she is the mother of a large family of children all of which is the most unusual and healthy to all external appearance.

And I think had she continued to ride on horse back
the worse horse made the
slower woman would
even have more

177
Surgery December 9th 1836

Case 2nd The case of Lady Shoemaker in this City living betwixt
Arche and Race in 11th St. This man had a most malignant
form of acquired scrofula - labouring under this disease
some time he had at length become quite much
emaciated the man had given up all hopes of a recov-
ery - and was not until this time that I was called
in I advised him to give up business and take free
exercise in the open air and after long persua-
sion I succeeded in accomplishing my request
the man through his industry and close attention
to business had acquired a little fortune - he pur-
chased a farm in the State of Pennsylvania and
removed from the City to his farm in
County - and had the pleasure of seeing
him in the street 2 or 3 weeks past and to my
surprise & astonishment he has become
the most corpulency - imaginable and whose
cheeks are the very index of health
I would merely state to you that it had been about
three years since he left City - and he in-
formed me out of the 365 days in the year he
spent 300 - at work on his farm or some
other active employment in the open air
never allowing himself to stay in the house
only at such times as was absolutely neces-
sary - such as meal times - and the house
for sleep.

Those 2 Cases show the importance of free exercise
in the open air - in the disease of scrofula - With

Dec^r 12th 1836

J. Wiley

Martin L. Symonds

Treat of acquired scrofula

Diet should consist of vegetables such as state bread crackers or biscuits with a
little butter, Milk, weak coffee for breakfast - For dinner may use in addition potatoes
turnips &c. I recommend you to Dr Colburn in diet for further particulars G.M.

Treatment of Acquired Scrofula Dec 13th 1836

Free exercise in open air, warm clothing, changing the dress every night & wearing flannel continually. Frictions with salt & water over whole surface of body every morning or morning & evening. Laxative & alterative medicines. Those laxatives should be used that stimulate the intestines to the evacuation merely of the feces. Blue pill also has a good effect on liver.

Preparations of iron They are the best tonics that can be used. Some use the carbonate others the tartrate & generally use the sulphate of iron.

Sedative It is a powerful stimulant tonic useful only in those states of the system connected with debility or atony.

After condition of stomach & bowels are regulated the diet must be improved. Allow the patient animal food especially that of white meat fowls.

Genuine cancerous tumor never occurs in corpulent persons.

Injuries & diseases of Head

Slight injuries of scalp should be attended to even burns as the inflammation is apt to extend to contiguous structures.

To these trivial injuries apply mild applications avoid exposure. Take no stimulating diet nor drinks use some mild purgative.

When coma or delirium supervenes & felt by convulsions no operation can relieve the patient. It is owing to a species of subacute inflammation or congestion in brain.

Active treat does not. Avoid general blood letting. Draw it locally apply cold applications to head. Rubrics to neck warm applications to feet. May be necessary to make an incision down to diseased bone.

To confine edges of wounds either lacerated or incised do not use suture. But cleanse the surface remove the roots of hairs & confine the edges by means of hair. Over this apply felt or lint. Keep parts cool by means of spirits & water.

Structures of Bones of Skull

When there is an opening in ~~contiguity~~ ^{contiguity} there will be considerable irritation. When there is a slight fissure only the inflammation is likely to extend to brain.

Injuries of the Head Dec. 18th 1836

When the integ remain entire even when the skull is broken into several fragments the effect will seldom be severe

Dec. 14th Fractures of Cranium

We never think of performing the operation of trephining when there is merely a fracture. If the depression be not very irregular you should not divide the scalp to examine & replace them.

To prevent inflammation keep patient quiet & use antiphlogistic remedies. Cold applications as bladder containing water.

Avoid all kinds of irritations. Use laxatives every day.

In even depressions the patient will recover without inconvenience.

If however the scalp has been torn open & the fragments are irregular they may be raised.

Where edges of fragments have been tilted in even where scalp remains sound it will be proper to make an incision. A T like incision will ^{generally} answer.

Sometimes it is best to make a crucial incision. Should apply saw only to sound bone where the edge is projecting. Should not be applied to depressed bone. If fragments is stripped of pericranium & of dura matter on inside remove them. If either of them adhere it is a moot point whether they shall be removed or not.

If there is an irreg depression with fracture even when the person remains sensible & there is no opening of matter yet the operation is required.

When any inveterate injury of brain remains under appropriate treatment apply the trephine.

Dec 16th Concussion Compression inflammation

Patient falls down Cold pale pulse feeble respiration feeble insensible.

Concussion There is the greatest irreg. in the symptoms.

Patient lies Cold senseless torpid & insensible.

In compression diff. in as much there is an attempt at reaction. In 5 to 10 or 20 days we may have another

clap of symptoms as those of puerperal fever. The symptoms are so various that you cannot classify them under any of these heads.

Treat of Concussion

Diminution of excitement is a favourable circumstance as the vessels are very often ruptured & if not they would (the vessels) would be overdistended provided the blood continues to rush on the head. Therefore should not be anxious to excite the system immediately. ^{merely remove neck etc} If there is danger of the patient's dying in this state depress the head you can aid this by frictions to extremities with hairs Volatiles to nose &c

In extreme cases which I have only met with 2 or 3 times that I had to use more powerful means. In one of the cases I could scarcely feel pulse in carotids. I injected hot brandy or into section in groins. Stimulating applications to skin as soon as you have succeeded in exciting the system be satisfied. When patient has so far recovered as to be able to swallow stimulents must not then be exhibited. Give water merely. If heat increases raise the head. Use moderately cold applics. Open bowels when patient can swallow.

And when reactions take place there is the time to bleed. Sometimes I have opened a vein & done in course of 10th

Give sedative powder as refrigerant & to help bowels open.

If injury occurs shortly after taking a meal give an emetic. Dec 20th Concussion does not require an operation.

Compression of Brain

Either from depression of bone, effusion of blood. There is a reaction in the arterial system. It manifests diff according to the part compressed. Paralysis pupils staring wide open. When they are compressed.

They require the unloading the vessels. Cold applications to head and other antiphlogistic treat.

Cases in which surgery is applicable. Where there is increasing depression &c when the sym cannot be allayed by antiphlog. It is not safe to operate in cases of depression when sym can be greatly relieved by appropriate treatment. Depletion general and local to as great an extent as patient can bear.

When compression exists on one side of cerebrum the paralysis will be on opposite side.

January 5th 1837 Chronic enlargement of tonsils

Chronic enlargement of scrophulous character only requires the operation

Foreign Bodies in Throat Run the finger boldly
down the throat & feel about The pharynx will be raised
By this means I can relieve 29 out of 30 patients
Needleman if patient vomits keep your finger down if you can
feel it

Cancer powder

Take of Litharge plaster 1 part
of Basilicon ointment 2 pts } 1 oz add 10 gr
of white oxide of Arsenic & apply it in form
of plaster to cancer G. McCallan

Jan. 12th 1837 Penetrating Wounds of Chest
1st Those merely penetrating into cavity
of pleura When adhesions do not exist between the
surfaces of pleura the lung will collapse & air will be drawn
into cavity of pleura. When full the air will be forced out
at every expiration & if the wound in the integuments be irregular
or oblique emphysema will take place into the ~~subcutaneous~~ cellular
tissue Should close the external wound by adhesive
plaster & over it a bandage The air will soon be absorbed
if not there may be made a small puncture at each angle
formed by the adhesive plaster &c The effusion of air
into cellular substance seldom becomes great in these cases
When the lungs are wounded Emphysema takes place into the
substance of the lungs and into anterior mediastinum & then over
the whole body & sometimes becomes enormously distended in this case
Close external wound apply bandage to chest Make small
punctures through integuments in diff parts If blood is effused
into cavity of pleura Search for the wounded artery if it be intercostal
or internal mannerly &c & close the wound so that it may unite
by 1st intention when blood will be absorbed
Bleeding purgatives &c will be good assistants

Medicines			Doses			Official prep	Dose
Emetics			Gr	3	3		
Specac.		1				Pulvis Specac. compositus & vinum Specac.	
Sulphas zinci		1				Maybe give in a tumbler of water	
Tartrate of Anti.	$\frac{1}{2}$ to $\frac{3}{4}$					Every 10 or 15 m. until it vomits in a tumbler	
Tobacco	10	+				Lincture dose is	$\frac{1}{2}$ to $\frac{3}{4}$
Lobelia In.	10	1				Dose of lincture	$\frac{1}{2}$ to $\frac{3}{4}$
Blue vitrol	5 to 15					Lincture of galls is its antidote	$\frac{1}{2}$ to $\frac{3}{4}$
Purgatives							
Elateum	$\frac{1}{2}$ to $\frac{1}{4}$					It may be given rubed up with sugar	
Elateum	$\frac{1}{2}$ to $\frac{1}{4}$						
Croton Tiglium	$\frac{1}{2}$ to 1 gtt.					It may be given in mucilage of gum Arabic	
Euphorbia Lathyrig	5 to 10 gtt.					Dose of oil It may be given rubed up with sugar	
Colocy. rhizidis	5 to $\frac{1}{2}$					Dose of Comp Ex Colocynth	
Black Hellebore	1	$\frac{1}{2}$				Dose of Bacher's tonic pills	5 gr. 3 to 4
Aloes	10					Lincture dose	$\frac{1}{2}$ to $\frac{1}{4}$
Scammony	3 to 5					It may be given triturated with sulphate of Potash sugar	
Tedachium Celt.	1	$\frac{1}{2}$					
All the above Cathartics are drastic in their operation							
Active cathartics							
Salap	1	$\frac{1}{2}$					
Rhubarb	1 to 2						
Scirra		1				Give in infusion	
Cassia Merilandica		1					
Juglans Cathartica	15 to $\frac{1}{2}$					of extract of inner bark	
Cassia Fistula		1				of extract	
Tamarind		1					
Magnesia	1	1					
Manna		1				to children	
Blue Pill	10 to 15					Alterative dose 4 or 5 gr	
John Wiley							
Antidote of hydrate of Soda							
Arsenic							

199

187


1881

Acacia gum
 Aqua Ammonica
 Rose Water
 Iode Pot-Carb
 Tartaric Acid
 Silver Shaped Bed pan
 C.C. Pills
 Alcohol
 Syr. Rhein aromat
 Chlorate of Potash
 Mt Potash
 1 lb. Lactate of Potash
 1 lb. Citron Oint
 Amices
 1 lb. Pot. Carb. Potash
 1 lb. Lig. Iodine Chlorinate
 43 Carb. Ammonium
 23 Precip. Pills
 43 Jalap. Pills
 Columbo root
 1 lb. Gentian root
 Bed pan bel. shaped
 2 Ergot Pills
 23 Opia Pills
 43 Acetas Plumb

1881

1 lb. Sarsaparilla Gum
 2 lb. Aqua Ammoniac
 1 qt. Rose Water
 3 lb. Soda Poi-Carb
 1 lb. Tartaric Acid
 1. Silver Shaped Bed pan
 100 C.C. Pills
 2 qt. Alcohol
 1 lb. Syr. Rhein aromat
 13 Chlorate of Potash
 1 lb. Nit Potash
 1/2 lb. Iodide of Potash
 1/2 lb. Citric Acid
 Amices
 1/2 lb. Bi. Carb. Potash
 1 qt. Lig. Sodae Chlorinate
 43 Carb. Ammonium
 23 Opie Pills
 43 Jalap Pulv
 Colombo root
 1 lb. Gentian root
 Bed pan bel. shaped
 2 Englat Pills
 23 Opie Pills
 43 Acetas Plumbe

1881





Transfused into Marshall Hall on Blood letting

on constitutional

JOHN

Detention of labor

causes 71 remedies

to relieve the patient

in various cases

General prep of each medicine

- | | | |
|---|-------------------|------|
| 1 | Salutarin of ant | 1 22 |
| 2 | Salutarin of ant | 30 |
| 3 | Hypo Nitrous acid | 38 |
| 4 | Albous acid | 46 |
| 5 | Nitric acid | 54 |

202
 $\frac{40}{3} \frac{1}{3} \frac{1}{2}$

1 = part in
 2 of one out

XXV

$\frac{24}{14}$
 38

How Powder

14 1/2

1 or 2

Rip

Div Powder

- 1 gr of opie
 1 gr of opie
 3 gr of Cal

To be given on going to bed

R

- | | |
|-----------------|-----|
| Muriate Ammonia | 3 1 |
| Acetat opii | 3 1 |
| Syrup Myrt | 3 1 |
| Alut. Liquorice | 3 6 |

Take desert spoon full
 3 W. daily

Samuel Jackson

R

- | | |
|----------------|---------|
| Alut. Tindus | } 3 1/2 |
| Hydriod. Ferri | |

Take 10 drops daily 3 t

in sweetened water R.J.

XXV



Distribution of 5th nerve
1st or ophthalmic division
It is purely nerve of sense
It divides into 3 branches
just as it is entering
through the foramen
lacrimum orbitale viz
internal which runs
to inner side of orbit
It sends off a branch
which forms with ~~the~~
Nasal division of 1st
branch of 5th pair the
Lacrimal ganglion
which lies on outside
of optic nerve

1837 1837 1837
Internal is felt on
It sends a nasal branch
to nose Other division
is infraorbital nerve
which runs out at inner
canthus of eye & ramifies
Frontal division of 1st
branch of 5th pair
3rd Lacrymal branch
2nd Branch of
5th nerve
Ganglion of Meckel
passes through sphenoid
maxillary fissure

Branches of Ganglion
of Meckel & 1st
ascending branch which
joins 2nd branch of 5th pair
branches which enter
nose = nasal branches
~~the maxillary nerve~~
& go through sphenoid
palatine foramen
one branch goes to
palate through posterior
palatine foramen
Median nerve goes off
from ganglion of
Meckel = Portia


Oculum Nerve describes
a retrograde curve & enters
introduces itself into
cranium through foramen
lacerum anterius
runs up on anterior
retrograde surface
to hyattus falopii
enters it which goes
into aqueductus
fallopia which contains
portio dura
& continues in connection
with it to
It finally joins the
ganglion of maxillary
nerve

Return to 2nd branch
of 5th pair
Where it pierces spheno
maxillary fissure
it sends 2 super
mular nerve
It then gets into
infra orbital fissure
making its appearance
on face through
infra orbital canal
2nd branch of 5th nerve
It pierces through foramen
ovale immediately
divides into 2 great bran
ches one

Wm Lloyd Garrison
New York

Wm

L.N.D. 3



Green's Chemistry

Nov^r 18th 1834

1834-11-9

1st primary form is has 6 sides

Prism has a flat surface at both ends

Light is similar to caloric in many of its prop^s Light of 2 kinds natural & artificial
Electricity When bodies are pos. excited

Electricity may be excited in all solid substances by friction Conductors & non
conductors Conductors = metals, charcoal, plumbago, water &c Non Conductors
belong glass, resins, sulphur, diamonds, dried wood, precious stones, silk, hair & wool
Part 3 Of ~~Considerable~~ bodies all substances are either electro positive or
electro negative There are but 5 simple electro-negative substances known
oxygen, chlorine, iodine, bromine & fluorine

